

Appendix H

Item 260-4: Handbook 133, Seed Count for Agriculture Seed

(Section 4.11. Procedure for Checking the Contents of Specific Agriculture Seed Packages
Labeled by Count)

Table of Contents

	Page
American Seed Trade Association (ASTA)/November 30, 2010.....	L&R - H3
AOSA Rules for Testing Seeds	L&R - H6
NIST Handbook 133, New Procedure for Testing Seed Count, A Call for Repeal Action Taken at the 95 th NCWM/January 24, 2011.....	L&R - H12
Association of American Seed Control Officials (AASCO)/January 5, 2010.....	L&R - H26

THIS PAGE INTENTIONALLY LEFT BLANK

AMERICAN SEED TRADE ASSOCIATION



November 30, 2010

John Gaccione
Chairman, NCWM Laws & Regulations Committee
c/o Westchester County Dep't of Weights & Measures
112 East Post Road, 4th floor
White Plains, NY 10601

Re: Seed Count Rule for Agricultural Seeds

Dear Mr. Gaccione:

The American Seed Trade Association (ASTA) is writing to express opposition to the pending petition from the Western Weights and Measures Association to rescind the seed count rule for agricultural seeds in Handbook 133 that was adopted by the National Conference of Weights and Measures (NCWM) in July 2010. We are deeply concerned by this proposal because of the uncertainty it will cause for our members and the inconsistency it will create for the regulatory community. The seed count rule adopted by NCWM is a validated, consistent, and practical means of verifying labeled seed counts for agricultural seeds. It would be detrimental to regulators and farmers if the rescission petition moves forward.

Benefits of the Seed Count Rule

At its annual meeting in 2010, the Conference adopted a rule regarding agricultural seed count that harmonized Handbook 133 with the provisions for verifying seed count that are used throughout the seed regulatory community. This action gave regulators an important tool to ensure the accuracy of labeled net quantity of contents statements on large bags of agricultural seed, which increasingly include a statement of count. This was particularly important for regulators in agricultural states because, prior to adoption of the seed count rule, there was no effective means for weights and measures regulators to verify labeled statements of count.

The adopted procedures incorporated the seed count rules that the Association for Official Seed Analysts (AOSA) developed as a result of many years of careful study and consideration. AOSA is a professional organization of seed laboratories whose members include state and federal seed regulators. Because AOSA's rules are used by seed regulators when they verify labeled statements of count, NCWM's action created uniformity in the laws for determining seed count for corn, soybean, field bean and wheat seed. Without this uniformity, certain states would apply different procedures and MAVs depending on whether the seed regulators or weights and measures inspectors were verifying the accuracy of seed count labeling.

225 Reinekers Lane • Suite 650 • Alexandria, VA 22314-2875 • (703) 837-8140 • Fax: (703) 837-9365

\\DC - 059737\000019 - 3172235 v1



The adopted procedures provide benefits to weights and measures regulators, farmers, and seed companies. Weights and measures regulators now have an effective and practical method to verify labeled seed count. This increases their ability to regulate labeling of these important commodities effectively. Farmers now have assurance that the seed they buy is accurately labeled. Seed companies now have assurance that a uniform testing procedure is in place for both weights and measures and seed regulators.

Concerns with the Rescission Proposal

The pending proposal would remove the seed count rule from Handbook 133 and establish a working group to consider next steps. Specifically, the working group would be directed to undertake “necessary studies, laboratory testing, field trial, and other appropriate measures to establish procedures for the verification of the accuracy and repeatability of ‘mechanical seed counter’ devices and/or to develop seed count procedures that are practical and reliable for field enforcement activities by Weights and Measures officials.” These steps are unnecessary and would be duplicative of the validation efforts already conducted by AOSA when the seed count rule was initially promulgated. Indeed, it is our understanding that the Conference originally rejected proposals to address this issue more than a decade ago due to the absence of the validating research that was subsequently undertaken by AOSA.

AOSA’s method verifies seed count with a mechanical seed counter. This method is widely used to verify seed count and was validated by the seed regulatory community. The rule is practical, straight forward, reliable, replicable, and cost effective. AOSA plans to submit to the Conference a white paper regarding the validation of its method. It is important to recognize that the procedure does not require any specialized knowledge or expertise to apply. Although its wording may sound technical, implementation is not. For example, the “purity analysis” in the rule simply requires separation of seeds from any debris such as sticks, rocks, and bugs that may be in the sample.

Although there may be concerns about the cost of mechanical seed counters, this does not justify rescission of the rule. There is no mandate to buy such equipment and, for states that do not expect to conduct inspections of seed count (i.e., non-agricultural states), it would not be worthwhile to do so. For states that do expect to do such inspections, purchase of seed counters will be a resource allocation issue that must be determined on a jurisdiction-by-jurisdiction basis. It is important to note, however, that most seed regulators and state seed laboratories already have mechanical seed counters that could be shared with their weights and measures colleagues. Furthermore, the Conference voted to adopt the seed count rule after discussing and debating cost concerns.

If the Conference adopts this proposal, it will rescind an efficient, practical, and effective means that weights and measures regulators have to verify seed count. Because the rule currently in place is validated and effective, efforts from a working group would be unnecessarily duplicative of AOSA’s previous work.

November 30, 2010
Seed Count Rule for Agricultural Seeds
Page 3



Implementation Assistance

Many of the concerns in the rescission proposal are appropriately addressed through training programs, as all new procedures inherently require training to ensure effective implementation. AOSA has offered to work with the Conference or interested states to disseminate training materials and also would welcome the opportunity to hold training webinars for interested states or regulators. Although no specialized knowledge is required to implement the seed count rule, such training programs will ensure that regulators have first hand knowledge of how the procedure works. Training, not amendments to Handbook 133, provide the traditional and appropriate vehicle for addressing implementation concerns.

In conclusion, ASTA opposes the petition to rescind the seed count rule because it would be a step backwards for the Conference. Keeping the rule in place is in the best interest of weights and measures regulators, seed regulators, farmers, and seed manufacturers.

If we may be of assistance or you have any questions, please do not hesitate to contact us. Thank you for your consideration.

Sincerely,

A handwritten signature in black ink that reads "A. W. LaVigne". The signature is written in a cursive, flowing style.

Andrew W. LaVigne
President & CEO
American Seed Trade Association

cc: Lisa Warfield, NIST
Don Onwiler, NCWM

AOSA Rules for Testing

- (1) Coated or encrusted seed: seed that has been covered by a layer(s) of materials that obscure the original shape and size of the seed resulting in a substantial weight increase. The addition of biologicals, pesticides, identifying colorants or dyes, and/or other active ingredients including polymers can be included in this process. Refer to sections 3.8 and 6.8 I.
- (2) Film-coated seed: film-coated seed retains the shape and the general size of the raw seed with a minimal weight gain. The film coating may contain polymers, pesticides, biologicals, identifying colorants or dyes, and other additives. The coating should result in a more or less continuous covering that eliminates or minimizes product dust-off.
- (3) Inoculated seed: seed that has received a coating of a commercial preparation containing a microbial product, e.g. *Rhizobium* sp.
- (4) Pelleted seed: seed that has been covered by a layer(s) of materials that obscure the original shape and size of the seed resulting in a substantial weight increase and improved plantability or singulation. The addition of biologicals, pesticides, identifying colorants or dyes, and/or other active ingredients including polymers can be included in this process. Refer to sections 3.8 and 6.81.
- (5) Raw seed: seed that is free of any applied materials.
- (6) Treated seed: seed with a minimal covering of various materials whose primary objective is to reduce or control certain disease organisms, insects or other pests attacking the seed or seedlings growing therefrom and that contains identifying colorants or dyes.

22 Obtaining the working sample

The working sample on which the actual analysis is performed shall be taken from the submitted sample in such a manner that it will be representative. A suitable type of mechanical divider (conical, centrifugal, riffle, etc.) should be used. To avoid damage when dividing large-seeded crop kinds such as beans, peas, etc., prevent the seeds from falling great distances onto hard surfaces. When dividing coated, encrusted, and pelleted seeds, mechanical dividers may be used only if the distance of the fall does not damage the applied materials.

For seed moisture determination, sub-samples must be drawn quickly to avoid exposing the seeds to the ambient air. Mechanical dividers are not appropriate for this purpose. Refer to section 2.2 b (3).

- a. Mechanical dividers. — This method is suitable for most kinds of seeds. The apparatus divides a sample into two approximately equal parts. The submitted sample is mixed by passing it through the divider, recombining the two parts and passing the whole sample through a second time and similarly a third time. After

mixing, the sample shall be reduced by passing the seed through the divider repeatedly, removing half the sample on each occasion. This process of successive halving is continued until a working sample of approximately, but not less than the minimum weight(s) stated in Table 2A is obtained.

Use of compressed air or a vacuum is highly recommended for cleaning mechanical dividers.

- (1) Centrifugal divider (Garnet type): This divider is suitable for all kinds of seed though it is not recommended for oilseeds (such as rapeseed, canola, mustards, flax) and kinds susceptible to damage (such as peas, soybeans, etc) and the extremely chaffy types.

The divider makes use of centrifugal force to mix and scatter seeds over the dividing surface. The seed flows downward through a hopper onto a shallow rubber cup or spinner. Upon rotation of the spinner by an electric motor the seeds are thrown out by centrifugal force and fall downward. The circle or area where the seeds fall is equally divided into two parts by a stationary baffle so that approximately half the seeds fall in one spout and half in the other spout. The centrifugal divider tends to give variable results when not carefully operated, and therefore the following procedure must be used:

- (a) Preparation of the apparatus:

- (i) Level the divider using the adjustable feet.
- (ii) Check the divider and four containers for cleanliness. Note that seeds can be trapped under the spinner and become a source of contamination.

- (b) Sample mixing:

- (i) Place a container under each spout.
- (ii) Feed the whole sample into the hopper; when filling the hopper, the seed must always be poured centrally.
- (iii) After the sample has been poured into the hopper, the spinner is operated and the seed passes into the two containers. Turn off spinner.
- (iv) Full containers are replaced by empty containers. The contents of the two full containers are fed centrally into the hopper together, the seed being allowed to blend as it flows in. The spinner is operated.
- (v) The sample mixing procedure is repeated at least once more.

- (c) Sample reduction:

- (i) Full containers are replaced by empty containers. The contents of one full container are set aside and the contents of the other container are fed into the hopper. The spinner is operated.
- (ii) The successive halving process is continued until the working sample(s) of not less than the minimum weight(s) required stated in Table 2A are obtained.

(iii) Ensure that the divider and containers are clean after each mixing operation.

(2) Soil/Riffle divider: This divider is suitable for most kinds of seed. For round-seeded kinds such as *Brassica* species, the collection containers should be covered to prevent the seeds from bouncing out.

This divider consists of a hopper with attached channels or ducts, a frame to hold the hopper, four collection containers and a pouring pan. Ducts or channels lead from the hopper to the collection containers, alternate ones leading to opposite sides. Riffle dividers are available in different sizes for different sizes of seed. The width and number of channels and spaces are important. The minimum width of the channels must be at least two times the largest diameter of the seed or any possible contaminants being mixed.

This apparatus, similar to the centrifugal divider, divides the sample into approximately equal parts.

(a) Preparation of the apparatus:

- (i) Place the riffle divider on a firm, level clean surface. Ensure the divider is level.
- (ii) Ensure that the divider and the four sample collection containers are clean. Check all channels, joints and seams of the divider and collection containers to ensure there are no seeds or other plant matter present before each use.
- (iii) Two clean empty collection containers shall be placed under the channels to receive the mixed seed.

(b) Sample mixing:

- (i) Pour the whole sample into the divider by running the seed backwards and forwards along the edge of the divider so that all the channels and spaces of the divider receive an equal amount of seed.
- (ii) The two full containers shall be replaced with two clean empty containers.
- (iii) The contents of one full container shall be poured into the divider by holding the long edge of the pan against the long edge of the riffle hopper and then rotating the bottom up so that the seeds pour across all channels at the same time, followed by the other full container using the same procedure.
- (iv) This process of mixing the entire submitted sample shall be repeated at least one more time before successive halving begins.

(c) Sample reduction:

- (i) The contents of one full container are set aside. Empty containers are placed under each channel, and the contents of the other container is poured into the hopper by holding the long edge of the pan against the

- (a) Preparation of the apparatus: Ensure that two trays, spatula and spoon are clean.
 - (b) Sample mixing:
 - (i) The sample is poured uniformly over a tray with a side to side swinging motion.
 - (ii) The receiving pan should be kept level.
 - (iii) This mixing procedure is repeated a minimum of three times.
 - (c) Sample reduction:
 - (i) A tray, a spatula and a spoon with a straight edge are required. After the preliminary mixing, pour the seed evenly over the tray with a side-to-side swing, alternately in one direction and at right angles to it. The depth of the seed in the pan shall not exceed the height of the vertical sides of the spoon. Do not shake the tray thereafter.
 - (ii) With the spoon in one hand, the spatula in the other, and using both, remove small portions of seed from not less than five random places on the tray.
 - (iii) Sufficient portions of seed are taken until the working sample(s) of not less than the minimum weight(s) required stated in Table 2A are obtained.
- (2) Hand-halving method: This method can be used when a proper mechanical divider is not available.

Procedure:

- (a) Seed is poured evenly onto a clean smooth surface.
 - (b) The sample shall be thoroughly mixed using a flat-edged spatula and placed into a pile.
 - (c) The pile shall be divided in half using a straight edge or ruler.
 - (d) Each half portion is divided in half.
 - (e) Each of the portions is divided into half again. There are now eight portions.
 - (f) Arrange the eight portions into two rows of four.
 - (g) Alternate portions should be combined to obtain two halves e.g. combine the first portion from row 1 with the second portion from row 2. Remove the remaining four portions.
 - (h) Repeat steps (a) to (g) until sufficient portions of seed are taken to constitute a working sample(s) of not less than the minimum weight(s) required stated in Table 2A are obtained.
- (3) For seed moisture determination, mix the submitted sample by tumbling or shaking the submitted sample bag, then open bag and use a spoon to remove portions from several random locations within the bag to obtain the appropriate working weight for one replicate. Place seeds in a moisture testing container. Repeat the procedure of mixing and sampling for the second replicate. Do not expose the sample to ambient air for more than one minute.

SECTION 12: MECHANICAL SEED COUNT

The following method shall be employed when using a mechanical seed counter to determine the number of seeds contained in a sample of soybean (*Glycine max*), corn (*Zea mays*), wheat (*Triticum aestivum*) and field bean (*Phaseolus vulgaris*).

12.1 Samples.

Samples for testing shall be of at least 500 grams for soybean, corn and field beans and 100 grams for wheat and received in moisture proof containers. Samples shall be retained in moisture proof containers until the weight of the sample prepared for purity analysis is recorded.

12.2 Seed counter calibration.

The seed counter shall be calibrated daily prior to use.

- (a) Prepare a calibration sample by counting 10 sets of 100 seeds. Visually examine each set to insure that it contains whole seeds. Combine the 10 sets of seeds to make a 1,000 seed calibration sample. The seeds of the calibration sample should be approximately the same size and shape as the seeds in a sample being tested. If the seeds in a sample being tested are noticeably different in size or shape from those in the calibration sample, prepare another calibration sample with seeds of the appropriate size and shape. Periodically re-examine the calibration samples to insure that no seeds have been lost or added.
- (b) Carefully pour the 1,000 seed calibration sample into the seed counter. Start the counter and run it until all the seeds have been counted. The seeds should not touch as they run through the counter. Record the number of seeds as displayed on the counter read out. The seed count should not vary more than ± 2 seeds from 1,000. If the count is not within this tolerance, clean the mirrors, adjust the feed rate and/or reading sensitivity. Rerun the calibration sample until it is within the ± 2 seed tolerance. If the seed counter continues to fail the calibration procedure and the calibration sample has been checked to ensure that it contains 1,000 seeds, do not use the counter until it has been repaired.

12.3 Sample preparation.

Immediately after opening the moisture proof container, mix and divide the submitted sample, in accordance with section 2.2, to obtain a sample for purity analysis and record the weight of this sample in grams to the appropriate number of decimal places (refer to section 2.3 a). Conduct the purity analysis to obtain pure seed for the seed count test.

RULES FOR TESTING SEEDS

12.4 Conducting the test.

After the seed counter has been calibrated, test the pure seed portion from the purity test and record the number of seeds in the sample.

12.5 Calculation of results.

Calculate the number of seeds per pound to the nearest whole number using the following formula:

$$\text{Number of seeds per pound} = \frac{453.6 \text{ g/lb} \times \text{no. of seeds counted in d.}}{\text{weight (g) of sample analyzed for purity}}$$

12.6 Tolerances for results from different laboratories.

Multiply the labeled seed count or first seed count test result by four percent for soybean samples, two percent for corn (round, flat or plateless) samples, five percent for field bean samples and three percent for wheat samples. Express the tolerance (the number of seeds) to the nearest whole number. Consider the results of two tests in tolerance if the difference, expressed as the number of seeds, is equal to or less than the tolerance.

Example:

Kind of seed: Corn
Label claim (1st test): 2275 seed/lb.

Lab Test (2nd test): Purity working weight = 500.3 g
Seed count of pure seed = 2479 seeds

$$\text{Number of seeds per pound} = \frac{453.6 \text{ glib} \times 2479 \text{ seeds}}{500.3 \text{ g}} = 2247.6 \text{ seeds/lb}$$

Rounded to the nearest whole number = 2248 seeds/lb

Calculate tolerance value for corn:

multiply label claim by 2%
 $2275 \text{ seeds/lb} \times 0.02 = 45.5 \text{ seeds/lb}$;
rounded to the nearest whole number = 46 seeds/lb

Determine the difference between label claim and lab test:

$$2275 \text{ seeds/lb} - 2248 \text{ seeds/lb} = 27 \text{ seeds/lb}$$

The difference between the lab test (2nd test) and the label claim (1st test) is less than the tolerance ($27 < 46$); therefore, the two results are in tolerance.

Slide 1

NIST HANDBOOK 133
New Procedure for Testing Seed Count

WWMA Call for Repeal
of
Action Taken at 95th Annual Meeting of
National Conference on Weights & Measures
Re:
Testing Procedures for Seed Count


Presented January 24, 2011, by
Kurt Floren
Director of Weights & Measures
County of Los Angeles, California



1

Slide 2

Association of Official Seed Analysts
“Rules for Testing Seeds”



AOSA: Official nationwide association of seed analysts, formed in 1908 in response to actions by individual states to develop seed laws.

Members include state, federal & university seed labs of U.S. and Canada.

Primary Functions:

- Establish AOSA Rules for Testing Seeds, adopted by most states as seed rules
- Contribute to refinement & modification of rules and procedures for seed testing
- Ensure procedures are standardized between analysts and between labs
- Influence and assist in enforcement of appropriate seed legislation at state and federal levels

2

Slide 3

Procedure does NOT start with simple count of 10 groups of 100

AOSA Rules for Testing Seeds
Volume 1. Principles and Procedures

SECTION 1: SAMPLING

Seed testing begins with the sampling process. No matter how accurately an analysis is made, it can show only the quality of the sample submitted; therefore, it is the responsibility of the seed sampler to insure that the sample is representative of the seed lot.

1.1 General procedure

- a. To secure a representative sample, equal portions shall be taken from evenly distributed parts of the quantity of seed to be sampled. Access shall be available to all parts of that quantity.
- b. For free-flowing seed in containers or bulk, a probe or trier long enough to sample all portions of the container shall be used. The trier should be able to remove an equal volume of seed from each part of the container through which the trier travels. Partitioned triers should be used when sampling vertically in the containers. Non-partitioned triers must be inserted into the containers diagonally or horizontally and should never be used when sampling vertically from the containers.
- c. Non free-flowing seed, such as chaffy grasses, uncleaned seed, large fragile seed, screenings, and seed with a low moisture content, that are difficult to sample with a probe or trier can be sampled by the hand sampling method. When a sample is taken by hand, insert the hand straight with the fingers together. Keep the fingers together as the hand is closed and withdrawn. Hand samples should be taken from various locations in the containers. Additionally, coated seed, seed mats, and seed tapes can be sampled by hand.
- d. Each probe, trier, or handful of seed removed from the lot is considered a primary sample. Each primary sample should be visually checked for uniformity. When the primary samples appear to be uniform, they shall be combined to form the composite sample prior to submission for testing. If non-uniformity is observed in the primary samples, the sampler should either discontinue sampling and advise management of the findings or continue sampling being sure to record the observations of non-uniformity on the sampling documentation.

1.2 Sampling equipment

The sampler must determine the most appropriate sampling tool and technique for the particular sampling situation. Manual sampling tools should be able to reach all portions of the container and have openings at least two and one half times the maximum diameter of the seed and possible contaminants that could be within the lot being sampled. Types of triers which can be utilized are the Noble type, open tube single sleeve, open tube double sleeve, non-compartmentalized double sleeve, compartmentalized double sleeve, and the pelican type sampler.

1-1
2009

To secure a representative sample, **equal portions** shall be taken from **evenly distributed parts**....

...a **probe or trier**...shall be used... able to remove an equal volume... from each part of the container...

Each probe, trier, or handful... is considered a primary sample. Each should be visually checked for uniformity.

...determine the most appropriate tool and technique...Manual sampling tools should be able to reach all portions of the container and have openings at least 2½ times the maximum diameter of... seed and possible contaminants...

Slide 4

Procedure does NOT start with simple count of 10 groups of 100

AOSA Rules for Testing Seeds

Note: The trier commonly referred to as the "Third Trier" is not appropriate for obtaining a representative seed sample, because it obtains seed only from the outermost portion of the container.

1.3 Obtaining a sample for submission

a. Seed in containers (typically packaged in 60 pounds or less)

- (1) When more than one probe or trier of seed is drawn from a container, follow different paths. When more than one handful is taken from a container, take them from well-separated points.
- (2) For lots of one to six containers, sample each container and take at least five primary samples.
- (3) For lots of more than six containers, sample five containers plus at least 10% of the number of containers in the lot. Round numbers with decimals to the nearest whole number. Regardless of the lot size, it is not necessary to sample more than 30 containers.

Number of containers in lot	Minimum number of containers to sample	Minimum number of containers in lot	Minimum number of containers to sample	Number of containers in lot	Minimum number of containers to sample
1-4	1 in 100 (one)	5-14	5	15-24	5
5	5	15-24	5	25-34	5
6	6	25-34	5	35-44	5
7-14	5	45-54	5	55-64	5
15-24	5	65-74	5	75-84	5
25-34	5	85-94	5	95-104	5
35-44	5	105-114	5	115-124	5
45-54	5	125-134	5	135-144	5
55-64	5	145-154	5	155-164	5
65-74	5	165-174	5	175-184	5
75-84	5	185-194	5	195-204	5
85-94	5	205-214	5	215-224	5
95-104	5	225-234	5	235-244	5
105-114	5	245-254	5	255-264	5
115-124	5	265-274	5	275-284	5
125-134	5	285-294	5	295-304	5

b. Seed in unit-bulk containers (typically containing 1,000 to 5,000 pounds)

Lot Size	Number of primary samples to be taken
1 Container	At least 3 primary samples from different sections of the container. At least 6 primary samples.
2 - 10 Containers	If fewer than 6 containers are in the lot, an equal number of primary samples must be taken from each container. At least 6 primary samples.
11 Containers or more	At least 6 primary samples, each sample drawn from a different container.

1-2
2009

For lots of one to six containers, sample each...
...take at least five primary samples

For lots of more than six containers, Sample five...plus at least 10% of the number of containers in the lot. (up to 30 primary samples)

Samples are drawn to form composite sample.

All of this is required for Purity Analysis

Procedure Requires Lab Analysis

AOSA Rules for Testing Seeds

(6) Two and three seed samples (as compared by AOSA Rules for Testing Seeds Vol. 3 Uniform Classification of Weed and Crop Seeds) shall consist of at least 600 seeds per sample for germination purposes (1,000 seeds for paired tests). If a purity analysis or a noxious-weed seed examination is required, the subsampled sample shall provide at least the minimum weights of working samples set forth in section 2.4.

(7) Coated, encrusted, or pelleted seed submitted for testing shall consist of at least: 7,500 seed units for a purity analysis, 30,000 seed units for a noxious-weed seed examination, or 1,000 seed units for a germination only test. Refer to sections 3.8 and 6.8.1.

b. **Individual-container samples being tested for uniformity**

The size of any individual container sample to determine uniformity in a seed lot shall be not less than the quantities indicated in the column "Minimum weight for noxious-weed seed or bulk examination" for the respective kinds of seed listed in Table 2A. If the sample drawn is larger than required, it shall be thoroughly mixed before it is divided to the desired size.

1.5 Forwarding of seed samples to laboratory

- a. After the appropriate number of primary samples for the size of the seed lot are drawn and combined into the composite sample, the entire composite sample is submitted to the laboratory. If the composite sample is too large, or if the composite sample is to be split into duplicate samples for submission to the same or different laboratories, the composite sample must be thoroughly blended and then split by an appropriate mechanical or non-mechanical method (refer to sec. 2.2).
- b. The person who samples the seed shall insure that the seed sample container is appropriately closed and sealed, if required, is identified in such a manner as to correlate with the sampling documentation, and is initialed and dated. Treated seed samples should be packaged in nonporous containers and be identified with the name of the applied seed treatment.
- c. Samples submitted for moisture testing or mechanical seed counts shall be packed in moisture-proof containers.
- d. Coated, encrusted, or pelleted seed shall be forwarded in firmly packed crush-proof, moisture-proof containers.
- e. Seeds are living organisms that can suffer damage in transit. Whenever possible, care should be taken to minimize damage during transit to the laboratory.

1 - 4
2009

After...appropriate number of primary samples...are drawn and combined into the composite sample,
the entire sample is submitted to the laboratory

As you will see, the procedure adopted into Hdbk 133 requires Purity Analysis testing by a seed analysis laboratory

5

Why be concerned with portions not adopted into Hdbk 133?

AOSA Rules for Testing Seeds
Volume 1. Principles and Procedures

SECTION 4: UNIFORM CLASSIFICATION OF WEED AND CROP SEEDS

For classification of weed and crop seeds, refer to AOSA Rules for Testing Seeds Volume 3, Uniform Classification of Weed and Crop Seeds, AOSA 2008, and subsequent updates. This entire handbook shall be considered part of the Rules and its use is required for determination of classification of the kind of seed under consideration and classification of weed and crop seed contaminants for purity testing.

4.1 Classification of all contaminating species shall be based on the species classification assigned to the kind or kinds under consideration. To determine the appropriate classification of a contaminating species:

- a. Under the name of each kind(s) under consideration determine the species classification, see column 4 (type class) of the table in the classification section of Volume 3. Species classification will be assigned to one or more of the following categories: agricultural (A), forest (F), herb or spore (H), nonparasitic and rampant (N), shrub and tree (S), turf (T), or vegetable (V).
- b. Under the name of the contaminating species find the classification (either weed or crop) in the corresponding column of contaminating classification based on the species classification (either A, F, H, R, S, T or V) for the kind under consideration. See page 4 of Volume 3 for an example.
- c. For exceptions to the classification system see items 2, 3 and 5, on page vi, of Volume 3.
- d. For guidelines on the classification of contaminating species found in the purity analysis of a single kind with multiple species classifications or to mixtures of kinds with different species classifications, see page vi, item 6, of Volume 3.

4 - 1
2009

This **entire handbook** shall be considered part of the Rules and **its use is required** for determination of classification of the kind of seed under consideration and classification of weed and crop seed contaminants for **purity testing**

Purity Testing is a requisite part of procedure adopted by NCWM

6

Slide 7

Seed Count Testing Procedure References AOSA “Rules” § 2.2

AOSA Rules for Testing Seeds
Volume 1. Principles and Procedures

SECTION 2: PREPARATION OF WORKING SAMPLES

The laboratory analysis for law enforcement, labeling, and general information as to seed quality, should determine the following for the sample analyzed: (1) the purity composition, (2) the rate of occurrence of noxious-weed seeds per unit weight, and (3) the percentage germination of the pure seed under consideration. Additional information, such as, seed count, detection of seed treatment, bulk examination for contaminants, tetrazolium visibility, detection of fungal endophytes, and seed moisture content may be determined using approved procedures.

2.1 Definitions:

- a. **Kind:** (of seed) one or more related species or subspecies that singly or collectively is usually known by one common name.
- b. **Seed unit:** the structure usually regarded as a seed in planting practices and in commercial channels. Refer to section 3.2 e for pure seed unit descriptions.
- c. **Working samples:**
 - (1) **Bulk examination working sample:** the sub-sample taken from the submitted sample on which a bulk examination is performed. Refer to sections 2.3 b and 3.3.
 - (2) **Fungal endophyte working sample:** the sub-sample of either seed or tillers taken from the submitted sample on which the fungal endophyte test is performed. Refer to sections 2.3 d and 10.
 - (3) **Germination working sample:** the sub-sample taken either from the pure seed portion of the purity analysis or directly from the submitted sample on which the germination test is performed. Refer to sections 2.3 c and 6.1.
 - (4) **Purity working sample:** the sub-sample taken from the submitted sample on which the purity analysis is performed. See sections 2.3 b and 3.
 - (5) **Noxious weed seed examination working sample:** the sub-sample taken from the submitted sample on which the noxious-weed seed examination is performed. Refer to sections 2.3 b and 3.1.
 - (6) **Seed moisture determination working sample:** the sub-sample taken from the submitted sample on which the seed moisture determination is performed. See sections 2.3 e and 11.
- d. **Seed format:**

2-1
2009

The laboratory analysis for **law enforcement, labeling...** should determine the following:

- (1) the **purity composition**
- (2) the rate of...noxious-weed seeds
- (3) the percentage germination...

By making reference to Section 2.2 (in Section 12 adopted by NCWM) all of Section 2 must be followed to ensure that seed count verification testing is defensible under legal challenge (i.e., defense in prosecution)

Slide 8

Seed Count Testing Procedure References AOSA “Rules” § 2.2

AOSA Rules for Testing Seeds

- (1) **Coated or encrusted seed:** seed that has been covered by a layer(s) of materials that obscure the original shape and size of the seed resulting in a substantial weight increase. The addition of biologicals, pesticides, identifying colorants or dyes, and/or other active ingredients including polymers can be included in this process. Refer to sections 3.8 and 6.8.1.
- (2) **Film coated seed:** film coated seed retains the shape and the general size of the raw seed with a minimal weight gain. The film coating may contain polymers, pesticides, biologicals, identifying colorants or dyes, and other additives. The coating should result in a more or less continuous covering that eliminates or minimizes product dust off.
- (3) **Inoculated seed:** seed that has received a coating of a commercial preparation containing a microbial product, e.g. *Rhizobium* sp.
- (4) **Pelleted seed:** seed that has been covered by a layer(s) of materials that obscure the original shape and size of the seed resulting in a substantial weight increase and improved plumbability or impaction. The addition of biologicals, pesticides, identifying colorants or dyes, and/or other active ingredients including polymers can be included in this process. Refer to sections 3.8 and 6.8.1.
- (5) **Raw seed:** seed that is free of any applied materials.
- (6) **Treated seed:** seed with a minimal covering of various materials whose primary objective is to reduce or control certain disease organisms, insects or other pests affecting the seed or seedlings growing therefrom and that contains identifying colorants or dyes.

2.2 Obtaining the working sample

The working sample on which the actual analysis is performed shall be taken from the submitted sample in such a manner that it will be representative. A suitable type of mechanical divider (conical, centrifugal, riffle, etc.) should be used. To avoid damage when dividing large-seeded crop kinds such as beans, peas, etc., prevent the seeds from falling great distances onto hard surfaces. When dividing coated, encrusted, and pelleted seeds, mechanical dividers may be used only if the distance of the fall does not damage the applied materials.

For seed moisture determination, sub-samples must be drawn quickly to avoid exposing the seeds to the ambient air. Mechanical dividers are not appropriate for this purpose. Refer to section 2.2 b (3).

- a. **Mechanical dividers.** – This method is suitable for most kinds of seeds. The apparatus divides a sample into two approximately equal parts. The submitted sample is passed by passing it through the divider, accumulating the two parts and passing the whole sample through a second time and similarly a third time. After

2-3
2009

The working sample... **shall be taken from the submitted sample**

“Submitted sample” means that submitted to the **seed laboratory** (recall Section 1.5)

A suitable type of **mechanical divider** (conical, centrifugal, riffle, etc.) **should be used**

Mechanical dividers are costly, sensitive pieces of equipment that Weights & Measures agencies do not possess

Slide 9

Examples of Mechanical Dividers



Dividing seed with mechanical divider

Preliminary research re: cost of Centrifugal Divider: \$2400 - \$3000

Centrifugal divider photos from AOSA presentation to IPSA



Dividing seed with mechanical divider




Riffle Divider: ~ \$400 - \$700



Boerner Divider: ~ \$1500 - \$1600

Slide 10

Purity Analysis: Only The Basics



...purity analysis...

determine the physical composition of the working sample.

The analysis shall include the identification of the kind... of seed under consideration, and all contaminating species and inert matter

The purity working sample shall be separated into the following components:

- (1) kind or cultivar to be considered **pure seed**
- (2) other crop seed
- (3) inert matter, and
- (4) weed seed

Requires seed identification expertise not possessed by most W&M officials

Slide 11

Pure Seed Sample: Required for Count Testing

Volume 1. Principles and Procedures

Purity analysis:	Percent by weight
Canada wildrye	92.13
Other crop seed	2.97
Inert matter	4.35
Weed seed	0.85

(C) 1000-seed separation of Pima egg.

Purity analysis:	Percent by weight
Pure seed (Pima egg)	97.25
Other crop seed (buckeye, blonduum vulgaris)	0.82
Inert matter	2.73
Weed seed	0.00

Kind	Weight (g)	Percent by weight	Percent of sample
410 common vetch (Pisica sativa ssp. sativa)	21.61	40.80 ÷ 0.9715 =	48.43
168 winter vetch (Pisica sativa ssp. sativa)	22.72	20.00 ÷ 0.9715 =	48.43
2 purple vetch (Pisica benghalensis)	0.00	0.20 ÷ 0.9715 =	0.19
			97.05

The percentages of common and winter vetch are listed separately as pure seed components of a section. The percentages of buffer from the purity analysis and purple vetch from the 1,000-seed separation are added together and included under other crop.

Purity analysis:	Percent by weight
Common vetch	48.43
Winter vetch	48.43
Other crop seed	0.21
Inert matter	2.73
Weed seed	0.00

For 400, 800, or 1000 seed separation tolerances refer to section 13, Table 13H.

For determining the rate of occurrence of any noxious-weed species that are similar in appearance to the kind or cultivar under consideration, refer to section 3.1.

3.2 Kind or cultivar considered pure seed. The pure seed shall include all seed units of each kind or each kind and cultivar under consideration present in excess of five percent of the whole. Kinds or kinds and cultivars present to the extent of five percent or less of the whole may be considered pure seed if shown on a label as components of a mixture in amounts of five percent or less.

3.3
2009

The **pure seed** shall include all seed units of each kind or each kind and cultivar under consideration...

Identification / determination of a PURE SEED sample is critical to the procedure and to demonstration of compliance with the Hdbk 133 procedure...

Are W&M field officials trained, qualified, certified?

11

Slide 12

Can't we just count 1,000 seeds?

AOSA Rules for Testing Seeds
 Volume 1. Principles and Procedures

SECTION 4: UNIFORM CLASSIFICATION OF WEED AND CROP SEEDS

For classification of weed and crop seeds, refer to AOSA Rules for Testing Seeds Volume 3 Uniform Classification of Weed and Crop Seeds, AOSA 2008, and subsequent updates. This entire handbook shall be considered part of the Rules and its use is required for determination of classification of the kind of seed under consideration and classification of weed and crop seed contaminants for purity testing.

4.1 Classification of all contaminating species shall be based on the species classification assigned to the kind or kinds under consideration. To determine the appropriate classification of a contaminating species:

- Under the name of each kind(s) under consideration determine the species classification; see column 4 (opp. class) of the table in the classification section of Volume 3. Species classification will be assigned to one or more of the following categories: grassland (G), flowers (F), herbs or spores (H), revegetation and rangeland (R), shrub and trees (S), leaf (L), or vegetable (V).
- Under the name of the contaminating species find the classification (either weed or crop) in the corresponding column of contaminating classification based on the species classification (either A, F, H, R, S, T or V) for the kind under consideration. See page v of Volume 3 for an example.
- For exceptions to the classification system see items 2, 3 and 5, on page vi, of Volume 3.
- For guidelines on the classification of contaminating species found in the purity analysis of a single kind with multiple species classifications or in mixtures of kinds with different species classifications, see page vi, item 6, of Volume 3.

4-1
2009

RECALL:

This **entire handbook** shall be considered **part of the Rules** and **its use is required** for determination of classification of the kind of seed under consideration and classification of weed and crop seed contaminants **for purity testing**

The procedure adopted by NCWM specifically requires calibration of mechanical seed counter using seed from a **Pure Seed** sample.....

Section 12.4 (Adopted in Hdbk 133):
"After the seed counter has been calibrated, test the pure seed portion from the purity test..."

12

Slide 13

Section 12 of “Rules for Testing Seeds” Directly Adopted in NCWM Action

AOSA Rules for Testing Seeds
Volume 1. Principles and Procedures

SECTION 12: MECHANICAL SEED COUNT

The following method shall be employed when using a mechanical seed counter to determine the number of seeds contained in a sample of soybean (*Glycine max*), corn (*Zea mays*), wheat (*Triticum aestivum*) and field bean (*Phaseolus vulgaris*).

12.1 Samples.

Samples for testing shall be of at least 100 grams for soybean, corn and field beans and 100 grams for wheat and received in moisture proof containers. Samples shall be retained in moisture proof containers until the weight of the sample prepared for purity analysis is recorded.

12.2 Seed counter calibration.

The seed counter shall be calibrated daily prior to use.

(a) Prepare a calibration sample by counting 10 sets of 100 seeds. Visually examine each set to insure that it contains whole seeds. Combine the 10 sets of seeds to make a 1,000 seed calibration sample. The seeds of the calibration sample should be approximately the same size and shape as the seeds in a sample being tested. If the seeds in a sample being tested are noticeably different in size or shape from those in the calibration sample, prepare another calibration sample with seeds of the appropriate size and shape. Periodically re-examine the calibration samples to insure that no seeds have been lost or added.

(b) Carefully pour the 1,000 seed calibration sample into the seed counter. Start the counter and run it until all the seeds have been counted. The seeds should not touch as they run through the counter. Record the number of seeds as displayed on the counter read out. The seed count should not vary more than ±2 seeds from 1,000. If the count is not within this tolerance, close the curtain, adjust the feed rate and/or reading sensitivity. Rerun the calibration sample until it is within the ±2 seed tolerance. If the seed counter continues to fail the calibration procedure and the calibration sample has been checked to insure that it contains 1,000 seeds, do not use the counter until it has been repaired.

12.3 Sample preparation.

Immediately after opening the moisture proof container, mix and divide the submitted sample, in accordance with section 2.2, to obtain a sample for purity analysis and record the weight of this sample in grams to the appropriate number of decimal places (refer to section 2.1.4). Conduct the purity analysis to obtain pure seed for the seed count test.

12-1
2009

The following method **shall be employed** when using a **mechanical seed counter**...

Samples shall be retained in moisture proof containers until the weight of the sample prepared for **purity analysis** is recorded.

...after opening the moisture proof container, **mix and divide** the **submitted sample**, in accordance with **section 2.2**, to obtain a sample for **purity analysis**

Conduct the purity analysis to obtain pure seed for the seed count test

Can there be any question that we are bound by the entire AOSA procedure?

13

Slide 14

Examples of “Pure Seed” Criteria

For Field Bean and Soybean:

- Seed with at least a portion of the seed coat attached
- Broken seed larger than one-half the original size with at least a portion of the seed coat attached
- For Fabaceae (*includes Field Bean & Soybean*):
 - Cotyledons that are broken apart but held together by the seed coat shall be classified as pure seed.
 - Cotyledons that have separated and are not held together by the seed coat are regarded as inert matter irrespective of whether or not the radicle-plumule axis and/or more than half of the seed coat may be attached.
- Wing, when present, is removed and considered inert matter.
- Chalcid-damaged seeds in Fabaceae that are puffy, soft, or dry and crumbly are considered inert matter.

Concern: Do typical Weights & Measures officials have such expertise?

14

Slide 15

Examples of “Pure Seed” Criteria


For Corn:

- Multiple floret, with or without awn, provided a caryopsis with some degree of endosperm development can be detected (either by slight pressure or by examination over light).
- Caryopsis or piece of broken caryopsis larger than one-half of the original size

Special Consideration:

- * A fertile floret attached to another fertile floret shall be separated
- * Attached glumes and empty florets shall be removed and classified as inert matter.

Concern: Again, does the average W&M official have such expertise?



15

Slide 16

Purity Analysis: Seed Identification

AOSA Rules for Testing Seeds
Volume 1. Principles and Procedures

SECTION 14: REPORT OF ANALYSIS

Laboratory reports of analysis that indicate laboratory testing was performed in accordance to the AOSA Rules for Testing Seeds are required to include, but not be limited to, the following information:

- Name and address of issuing laboratory.
- Name of responsible individual from the issuing laboratory.
- Unique laboratory test or sample number assigned by the issuing laboratory.
- Date report of analysis is issued.
- Applicant's information, such as kind of seed, cultivar, lot number, lot size, certification number, treatment, etc., as stated by the applicant.
- When the submitted sample is treated seed, inoculated seed, film-coated seed, coated or encrusted seed, or pelleted seed, this shall be indicated.
- Kind of seed by scientific name, or common name, or both.
- When a purity analysis is conducted the following information shall be reported under Purity Analysis:
 - Weight of purity working sample.
 - Percentage by weight of pure seed, other crop seed, inert matter and weed seed found in the purity working sample, given to two decimal places.
 - Scientific name, or common name, or both, of all other crop seed or weed seed found in the purity working sample. If none are found, this shall be indicated.
- When a noxious weed seed examination is conducted the following information shall be reported under Noxious Weed Seed Examination:
 - Weight of noxious weed seed working sample.
 - Scientific name, or common name, or both, of noxious weed seed found, the number of each type found and rate of occurrence per unit weight. If none are found, this shall be indicated.
- When a germination test is conducted the following information shall be reported under Germination Test:
 - Percentage of normal seedlings as a whole number (refer to section 6.7).
 - Percentage of hard seed, if applicable, as a whole number (refer to section 6.7).

14 - 1
2009

When a purity analysis is conducted, the following shall be reported under Purity Analysis:

- (1) Weight of purity working sample
- (2) Percentage...of pure seed, other crop seed, inert matter, and weed seed...
- (3) Scientific name, or common name, or both, of all other crop seed or weed seed found...

Seed Analysts typically work 4-5 years in a seed laboratory to gain expertise to independently conduct seed analyses...

What percentage of Weights & Measures officials are qualified? ANY?

16

Other Concerns: Repeatability?

AOSA Rules for Testing Seeds
Volume 1. Principles and Procedures

SECTION 12: MECHANICAL SEED COUNT

The following method shall be employed when using a mechanical seed counter to determine the number of seeds contained in a sample of soybean (*Glycine max*), corn (*Zea mays*), wheat (*Triticum aestivum*) and field bean (*Phaseolus vulgaris*).

12.1 Samples.

Samples for testing shall be of at least 500 grams for soybean, corn and field beans and 100 grams for wheat and received in moisture proof containers. Samples shall be retained in moisture proof containers until the weight of the sample prepared for purity analysis is recorded.

12.2 Seed counter calibration.

The seed counter shall be calibrated daily prior to use.

(a) Prepare a calibration sample by counting 10 sets of 100 seeds. Visually examine each set to insure that it contains whole seeds. Combine the 10 sets of seeds to make a 1,000 seed calibration sample. The seeds of the calibration sample should be approximately the same size and shape as the seeds in a sample being tested. If the seeds in a sample being tested are noticeably different in size or shape from those in the calibration sample, prepare another calibration sample with seeds of the appropriate size and shape. Periodically re-examine the calibration samples to insure that no seeds have been lost or added.

(b) Carefully pour the 1,000 seed calibration sample into the seed counter. Start the counter and run it until all the seeds have been counted. The seeds should not touch as they run through the counter. Record the number of seeds as displayed on the counter read out. The seed count should not vary more than ± 2 seeds from 1,000. If the count is not within this tolerance, clean the mirrors, adjust the feed rate and/or reading sensitivity. Rerun the calibration sample until it is within the ± 2 seed tolerance. If the seed counter continues to fail the calibration procedure and the calibration sample has been checked to insure that it contains 1,000 seeds, do not use the counter until it has been repaired.

12.3 Sample preparation.

Immediately after opening the moisture proof container, mix and divide the submitted sample, in accordance with section 7.2, to obtain a sample for purity analysis and record the weight of this sample in grams to the appropriate number of decimal places (refer to section 3.3 a). Conduct the purity analysis to obtain pure seed for the seed count test.

12-1
2009

Prepare a calibration sample by counting 10 sets of 100 seeds. Combine...to make a 1,000 seed calibration sample.

...pour...into the seed counter. ...run it until all seeds...counted. The seed count should not vary more than ± 2 seeds from 1,000.


If...not within this tolerance, clean...mirrors, adjust...feed rate and/or reading sensitivity. Rerun it until it is within the ± 2 seed count tolerance.

Calibration procedure mandates no steps to verify repeatability.
Out-Of-Tolerance runs could be unlimited.
Results may result in enforcement action: **Defensible?**

17

Mechanical Seed Counter

(Photo from AOSA presentation to IPSA)



Seen one? Own one? Have ready access to one?
Not subject to transportation on front seat of a pickup truck!
Preliminary Cost Estimate: \$8,000

18

Slide 19

Example of 100-Seed Sampling

(from AOSA presentation to IPISA)



Preparing calibration sample

Does this look like a field activity?

19

Slide 20

Other Concerns: Equipment Access? Portability?



Loading and running calibration sample.
(Photos from AOSA presentation to IPISA)



Use of mechanical seed counter is clearly **not a field operation**.

Equipment and analysis procedures are **laboratory activities**.

20

Slide 21

Adopted Hdbk 133 Procedure Does Not Duplicate AOSA Procedure

Key Difference:

- **AOSA** procedure sets **"TOLERANCES"**
[Corn: 2%; Wheat Seed: 3%; Soybean: 4%; Field Bean: 5%]
- **Hdbk 133** procedure sets **"MAVs"** [Same values]
(All references to "Tolerance" amended by L&R to "MAV" prior to adoption)

"Tolerance":

- Allows Underfill / Short Count in each package

"Maximum Allowable Variance" (MAV):

- Sets limit for any single Minus Error
- Average Error must still not be minus (>SEL)

21

Slide 22

Inspection Results on Same Lot Will NOT Result in Same Outcome

Key Questions:

- Does adopted procedure employ random sampling?
- Does adopted procedure permit action on LOT? (*Pass/Fail*)

Package Test Example (*Hypothetical*):

Lot Size: 200 50-lb. bags Corn (~80K seed ct. ea.)
Sample: 12 packages MAV/Tolerance: 2%
Errors: Minus 800 seeds (1%) in each sample
Avg. Error: - 1%

AOSA Procedure: Lot **PASSES** (No error > Tolerance)
Hdbk 133 Proc.: Lot **FAILS** (No UMEs, but Avg. Minus)

22

Slide 23

Call for Repeal of NCWM Adoption

Acknowledge:

1st: **We do need an accurate, reliable, uniform procedure for testing seed count.**

- Seed count is an important factor in farming to manage input costs & to meet needs of modern planting equipment
- Packers/Manufacturers are increasingly placing supplemental count statements on seed packages due to customer demand
- A procedure is needed by W&M to regulate labeled count accuracy

2nd: **AOSA standards are well established and are in wide use by seed labs.**

BUT: Procedure was prematurely adopted by NCWM.

- Procedure provides little assurance of counter accuracy.
- Procedure is not equivalent to AOSA (Tolerance vs MAV)
- We have adopted a **test procedure that few, if any, can actually perform!**

23

Slide 24

Call for Repeal of NCWM Adoption

NOT suggesting that procedure is not needed....

But,

New NIST Hdbk 133 sections 4.2 and 4.11

- Require expertise not held by W&M inspectors
- Require equipment not suitable for field use
- Require equipment that is cost restrictive
- Include procedures (e.g., "mix," "divide," "purity analysis," etc.) for which no guidance is given
- Require steps that, if not precisely followed, subject W&M agencies to legal challenges and, potentially, litigation exposure for taking off-sale action.

24


Slide 25

Recommendation

WWMA calls on NCWM to:

Recognize that:

- State & local W&M agencies **do not have required equipment**
- State & local W&M agencies **do not have required seed analysis expertise** (licensing/certification)
- State & local W&M agencies are highly unlikely to have time (years for certification as seed analysts) or resources (\$\$) to meet requirements
- Adopted procedures **do not facilitate field tests** of seed count
- Adopted procedures are **not equivalent** to AOSA method (Tol. vs MAV)
- Adopted procedures **will not result in enhanced enforcement** due to all of above.



25


Slide 26

Recommendation

WWMA calls on NCWM to:


Take the following actions:

- **Rescind action taken to adopt** amendments to Hdbk 133 Section 4.2 and to add Section 4.11 et seq
- Direct NCWM Laws & Regulations (L&R) Committee to *(at a minimum)*:
 - **Establish a working group** to conduct studies, field trials, laboratory testing, etc., to **establish procedures for verification of repeatability** of Mechanical Seed Counter devices
 - Revise proposed procedure to incorporate guidance to inspectors in conducting "mixing," "dividing," "purity analysis," and other steps
- Direct NCWM L&R Committee to **establish a working group to research, develop, and recommend alternative seed count testing procedures** that are practical and reliable for field applications *(preferred)*
- **VERY CAUTIOUSLY** consider the **ramifications of adopting any TOLERANCE** for any packaged commodity..... Slippery slope.



26

Slide 27





AASCO

Association of American Seed Control Officials

January 5, 2010

John Gaccione
Chairman, NCWM Laws & Regulations Committee
c/o Westchester County Dep't of Weights & Measures
112 East Post Road, 4th floor
White Plains, NY 10601

RE: Seed Count Rule for Agricultural Seeds

Dear Mr. Gaccione:

The Association of American Seed Control Officials (AASCO) has recently learned that there has been a petition from the Western Weights and Measures Association to rescind the seed count rule for agricultural seeds in Handbook 133. We respectfully urge you not to rescind this rule and we ask for your continued support of this rule as adopted by the National Conference of Weights and Measures (NCWM) in July 2010.

As stated last year in our letter to NCWM; Seed is a biological unit and as such, it is subject to environmental influences that introduce variation in size and density. Seed cannot be produced utilizing a standardized manufacturing process that controls size and density. Utilizing a process of referee testing and scientific review, The Association of Official Seed Analyst, has established in the Rules for Testing Seed, acceptable testing methods and variances for Corn, Soybeans, Field Beans, and Wheat that are recognized and utilized by state seed control officials in the administration of their respective state seed laws. These standards are also acceptable to the regulated seed industry as a fair and valid means for determining that their products are in compliance with the respective state seed laws.

The primary objective of the AASCO is to promote uniformity in the administration of seed laws. We feel that by rescinding this rule our efforts to promote uniformity between regulatory agencies would be greatly diminished. Further, with variations in seed count methodology being applied by seed regulatory officials, an unnecessary hardship will be created for the agriculture seed industry without tangible benefits to the end users. As such, the AASCO respectfully requests that existing rules used to establish seed counts and in harmony with rules established by the Association of Official Seed Analyst Rules for Testing Seed be left in place and not be rescinded.

Best Regards,

Ronald R. Pence
President – AASCO

Assistant Administrator, Commodity Inspection Division
Oregon Dept. of Agriculture
635 Capitol Street NE
Salem, Oregon 97301-2532

RONALD PENCE
President
Oregon Dept. Of Agriculture
Commodity Inspection Division
635 Capitol street N. E.
Salem, OR 97301-2532
Tel. 503-986-4620

JOHN HEATON
First Vice-President
Seed Services Director
Nursery, Cotton & Seed Services
1220 N Street, Room A-372
Sacramento, CA 95814
Tel. 916-654-0493

BRENDA BALL
Second Vice-President
Arizona Department of
Agriculture
1688 West Adams
Phoenix, AZ 86007
Tel. 602- 542-0986

LARRY NEES
Secretary
Office of Indiana
Seed Commissioner
175 S. University Street
Purdue University
West Lafayette, IN 47907-
2063
Tel. 765-494-1557

GREG HELMBRECHT
Treasurer
Wisconsin Department of Agr.,
Trade, & Consumer Protection
P. O. Box 8911
Madison, WI 53708-8911
Tel. 608-224-4596