



US010745753B2

(12) **United States Patent**  
**Almeida et al.**

(10) **Patent No.:** **US 10,745,753 B2**

(45) **Date of Patent:** **Aug. 18, 2020**

(54) **MOUSE CELL LINE AUTHENTICATION**

FOREIGN PATENT DOCUMENTS

(71) Applicant: **United States of America, as represented by the Secretary of Commerce, Gaithersburg, MD (US)**

WO WO/2016/066070 5/2016  
WO WO/2016/188144 12/2016

(72) Inventors: **Jamie Almeida, Eldersburg, MD (US); Kenneth Cole, Gaithersburg, MD (US)**

OTHER PUBLICATIONS

(73) Assignee: **Government of the United States of America, as Represented by the Secretary of Commerce, Gaithersburg, MD (US)**

Almeida et al. (BMC Biotechnology, 2011, vol. 11, p. 1-10, IDS reference) (Year: 2011).\*

Please see the attached listings on document 1: PRior Art for Information Disclosure Statement Submitted With Application-33 References) and document 2: (Prior art references from Cytotechnology paper).

(\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 442 days.

\* cited by examiner

(21) Appl. No.: **15/383,378**

*Primary Examiner* — Stephanie K Mummert

(74) *Attorney, Agent, or Firm* — Daphne Burton; Burton IP Law Group

(22) Filed: **Dec. 19, 2016**

(65) **Prior Publication Data**

US 2017/0101677 A1 Apr. 13, 2017

(57) **ABSTRACT**

**Related U.S. Application Data**

(63) Continuation-in-part of application No. 13/935,285, filed on Jul. 3, 2013, now Pat. No. 9,556,482.

A method, kit and primers for determining alleles present in a DNA sample that is sourced from either a mouse or a cell line derived from a mouse. A set of STR marker loci are selected that can be amplified together in a multiplex amplification reaction. The set of STR marker loci are selected from the group consisting of: 18-3, 4-2, 6-7, 15-3, 6-4, 12-1, 5-5, X-1, Mo11-1, Mo19-2, Mo1-2, Mo7-1, Mo1-1, Mo3-2, Mo8-1, Mo2-1, Mo11-2, Mo17-2, Mo13-1, D8S1106 and D4S2408. A set of oligonucleotide primer pairs are provided. Each oligonucleotide primer pair flanks a single locus in the set of STR marker loci, which are co-amplified in a multiplex amplification reaction, with the product(s) comprising a mixture of amplified alleles from each of the co-amplified loci in the set of STR marker loci. The products are evaluated to determine the alleles present at each of the loci analyzed.

(51) **Int. Cl.**  
**C12Q 1/68** (2018.01)  
**C12Q 1/6881** (2018.01)

(52) **U.S. Cl.**  
CPC ..... **C12Q 1/6881** (2013.01); **C12Q 2600/156** (2013.01); **C12Q 2600/16** (2013.01)

(58) **Field of Classification Search**  
None  
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

2013/0184176 A1 7/2013 Schumm

**33 Claims, 19 Drawing Sheets**

**Specification includes a Sequence Listing.**

STR MARKER	GENBANK ACCESSION #	LOCATION ON CHROMOSOME (BP)	PRIMERS (5'-3')	PRIMER $\mu$ M
18-3	NT_039674.8	60271556-60271705	F: [FAM] - <u>TC</u> TTCTCCTTTTGGTGCATGC R: <u>GTTTCTT</u> IGCTAAATAACTAAGCAAGTGAACAGA	0.200
4-2	NT_187032.1	82068280-82068580	F: [FAM] - AAGCTTCTCTGGCCATTTGA R: <u>GTT</u> CATAAACTTCAAGCAATGACA	0.125
6-7	NT_039353.8	51601265-51601685	F: [FAM] - AGTCCACCCAGTGCATTCCTC R: <u>GTTTCTT</u> ICATGTGGCTGGTATGCTGTT	0.075
9-2	NT_039474.8	74395400-74395000	F: [VIC] - GGATTGCCAAGAAATTTGAGG R: <u>GTTTCTT</u> TCCTGAGTTGTGGACAGGGTTA	0.080
15-3	NT_039617.8	4930200-4930500	F: [NED] - TCTGGGGTGTCTGTCAATA R: <u>GTTTCTT</u> TCTCAGGGAGGAGTGTGCT	0.060
6-4	NT_039360.8	142021975-142022270	F: [NED] - TTTGCAACAGCTCAGTTTCC R: <u>GTTTCTT</u> JAATCGCTGGCAGATCTTAGG	0.100
12-1	NT_039548.8	38480950-38481170	F: [VIC] - CAAAATTGTCATTGAACACATGTAA R: <u>GTTTCTT</u> CAATGGTCAAGAAAATACTGAAGTACAA	0.200
5-5	NT_109320.5	112641540-112641820	F: [PET] - CGTTTTACCTGGCTGACACA R: <u>GTTTCTT</u> TGGTTTAAAACTCAATACCAAACAA	0.300
X-1	NT_039706.8	110959842-110960080	F: [PET] - GGATGGATGGATGGATGAAA R: <u>GTTTCTT</u> AAGGTATATATCAAGATGGCATTATCA	0.300
D8S1106	NT_167187.1	12835860-12836150	F: [VIC] - GTTTACCCCTGCATCCTGG R: <u>GTTTCTT</u> TCAGAAATGCTCATAGTGCAAGA	0.150
D4S2408	NT_006316.16	31304210-31304514	F: [NED] - TCATTTCCATAGGGTAAGTAAAA R: <u>GTTTCTT</u> GCCATGGGGATAAAAATCAGA	0.200

FIG. 1

**SEQUENCING PRIMERS**

<b>STR MARKER</b>	<b>PRIMERS (5'-3')</b>	<b>AMPLICON SIZE (bp)</b>	<b>T(Celsius)</b>
18-3	18-3 F: TCTTTCTCCTTTTGGTGCATGC 18-3 R: <u>G</u> TCAAAGTTGGGGTTACAGAAATG*	281-313	54
4-2	4-2 F: AAGCTTCTCTGGCCATTTGA 4-2 R: <u>G</u> TTCATAAACTTCAAGCAATGACA	217-248	57
6-7	6-7 F: AGTCCACCCAGTGCATTCTC 6-7 R: <u>G</u> CATGTGGCTGGTATGCTGTT	333-515	60
9-2	9-2 F: GGCTCTCTCACACCTCATCC* 9-2 R: <u>G</u> TCCATGAATCCAGACATTCC	318-360	60
15-3	15-3 F: TCTGGCGTGTCTGTCATAA 15-3 R: <u>G</u> TTCTCAGGGAGGAGTGTGCT	157-222	60
6-4	6-4 F: TTTGCAACAGCTCAGTTTCC 6-4 R: <u>G</u> AATCGCTGGCAGATCTTAGG	276-311	52
12-1	12-1 F: CAAAATTGTCAATTGAACACATGTAA* 12-1 R: <u>G</u> CAATGGTCAAGAAAATACTGAAGTACAA	222-259	55
5-5	5-5 F: CGTTTTACCTGGCTGACACA 5-5 R: GATGCTTGCCTGTTCCCTAGC	258-298	60
X-1	X-1 F: GGATGGATGGATGGATGAAA X-1 R: <u>G</u> AAGGTAATATCAAGATGGCATTATCA	357-442	54

**FIG. 2**

**DEFINING STR FRAGMENT LENGTH WITH CORRELATING REPEAT NUMBER**

<b>STR</b>	<b>REPEAT MOTIF</b>	<b>ALLELE DISTRIBUTION AND CORRELATING FRAGMENT LENGTHS</b>
18-3	[ATCT] <sub>n</sub>	137 146 150 154 158 162 166 171 175 179 12.2 15 16 17 18 19 20 21 22 23
4-2	[GATA] <sub>n</sub> [GATG] <sub>n</sub> [ATAG] <sub>n</sub>	209 213 216 217 220 221 222 224 225 228 232 236 240 244 248 14 15 15.3 16 16.3 17 17.1 17.3 18 18.3 19.3 20.3 21.3 22.3 23.3
6-7	[CTAT] <sub>n</sub>	333 337 341 345 349 352 355 356 360 363 364 366 368 376 377 388 12 13 14 15 16 17 17.3 18 19 19.3 20 20.1 21 23 23.1 26
9-2	[TAGA] <sub>n</sub> [AGAT] <sub>n</sub>	188 192 204 219 220 223 227 231 235 7.1 8.1 11.1 15 15.1 16 17 18 19
15-3	[TAGA] <sub>n</sub>	157 165 169 177 181 184 189 193 196 200 204 208 212 216 220 222 12 14 15 17 18 18.3 19.3 20.3 21.3 22.3 23.3 24.3 25.3 26.3 27.3 28.1
6-4	[ATAG] <sub>n</sub> [ATGA] <sub>n</sub> [TAGA] <sub>n</sub>	276 285 286 289 290 293 294 297 298 301 302 307 311 11.2 13.3 14 14.3 15 15.3 16 16.3 17 17.3 18 19 20
12-1	[AGAT] <sub>n</sub> [GATA] <sub>n</sub>	222 226 227 230 231 234 235 238 239 242 246 247 250 254 259 15 16 16.1 17 17.1 18 18.1 19 19.1 20 21 21.1 22 23 24.1
5-5	[TATC] <sub>n</sub>	326 330 334 338 342 346 350 354 365 11 12 13 14 15 16 17 18 21
X-1	[ATAG] <sub>n</sub> [ATGA] <sub>n</sub> [TAGA] <sub>n</sub>	380 385 389 393 397 401 404 405 409 413 421 428 448 20 21 22 23 24 25 25.3 26 27 28 30 32 37

**FIG. 3**

**COMPLETE GENETIC PROFILES OF SIX MOUSE CELL LINES**

CELL LINE	ORIGIN	18-3	4-2	6-7	9-2	15-3	6-4	12-1	5-5	X-1
NIH3T3	NIH Swiss	17, 19	19.3, 19.3	12, 12	15, 16	20.3, 20.3	14.3, 14.3	20, 20	14, 15	25, 25
L-929	C3H/An	16, 16	20.3, 20.3	12, 12	15, 15	24.3, 25.3, 26.3	16, 16	16, 16	14, 14	26, 27
MC3T3-E1	C57BL/6	15, 15	20.3, 20.3	17, 17	17, 18	22.3, 22.3	17, 17	17, 17	17, 17	28, 28
RAW264.7	Balb/c	18, 18	22.3, 22.3	12, 12	15, 15	22.3, 22.3	17, 17	16, 16	14, 14	24, 24
P3X63Ag8.653	Balb/c	18, 19	21.3, 21.3	12, 12	15, 16	22.3, 23.3	17, 18	16, 16	13, 14	25, 25
HK-PEG-1	Balb/c	18, 19	21.3, 21.3	12, 12	15, 15	22.3, 23.3	17, 18	16, 16	13, 14	25, 25

**FIG. 4**




FIG. 5




FIG. 6





FIG. 7





MOUSE DNA SAMPLE	ORIGIN	MOUSE STR MARKERS										
		18-3	4-2	6-7	9-2	15-3	6-4	12-1	5-5	X-1		
CAST/EIJ	JL STOCK# 928	OUT OF RANGE	16	20	7.1	17	18	21.1	15	30		
MOLG/DN	JL STOCK# 555	20	17	23	7.1	17	18	19	14	32		
SWR/J	JL STOCK# 689	18	17.3	12	15	24.3	14.3	17	15	26		
129P3/J	JL STOCK# 690	18	19.3	12	16	20.3	17	17	14	26		
129S1/SVIMJ	JL STOCK# 2448	18	18.3	12	16	20.3	17	17	14	26		
BALB/CJ	JL STOCK# 651	17	21.3	12	15	21.3	17	16	13	25		
BUB/BNJ	JL STOCK# 653	17	19.3	12	15	20.3	16	18	15	26		
C3H/HEJ	JL STOCK# 659	16	19.3	12	15	26.3	18	16	15	27		
C57BL/6J	JL STOCK# 664	16	20.3	17	18	22.3	17	17	17	27		
C57L/J	JL STOCK# 668	16	18.3	15	15	18	17	17	17	26		
CBA/J	JL STOCK# 656	19	18.3	12	15	25.3	17	16	14	25		
DBA/1J	JL STOCK# 670	16	20.3	12	15	21.3	17	16	13	26		
I/LNJ	JL STOCK# 674	18	21.3	16	15	20.3	17	16	14	26		
KK/HLJ	JL STOCK# 2106	18	20.3	14	15	21.3	14.3	17	13	23		
LG/J	JL STOCK# 675	16	20.3	12	15	24.3	17	16	15	26		
LP/J	JL STOCK# 676	18	18.3	12	16	19.3	16.3	17	16	26		
LT/SVEI	JL STOCK# 3588	16	20.3	12	15	22.3	16	17	14	25		
NZB/BLNJ	JL STOCK# 684	22	19.3	18	15	21.3	15.3	18	12	24		
P/J	JL STOCK# 679	16	18.3	14	15	22.3	14.3	16	14	25		
PERC/EIJ	JL STOCK# 1307	18	19.3	20	17	21.3	16.3	18	14	25		

FIG. 9

MOUSE DNA SAMPLE	ORIGIN	MOUSE STR MARKERS									
		18-3	4-2	6-7	9-2	15-3	6-4	12-1	5-5	X-1	
SM/J	JL STOCK#687	20	18.3	12	15	25.3	14.3	16	14	26	
WSB/EJ	JL STOCK# 1145	19	20.3	14	16	18.3	17.3	16	14	21	
AJ	JL STOCK#646	18	21.3	12	15	22.3	17	16	16,17	27	
NOD/SHILTJ	JL STOCK# 1976	18	18.3	14	15	22.3	15.3	17	15	26	
NZO/HLLTJ	JL STOCK# 2105	19	19.3	19	15	22.3	16,17	17	12	24	
PWK/PH	JL STOCK# 3715	19	19.3	15	7.1	15	15	16.1,17.1	15	20	
CD1/ICR MALE	ZYAGEN#GM-150M	17	19.3	12,15	15	21.3	14.3,16	17,18	15	25	
CD1/ICR FEMALE	ZYAGEN#GM-150F	17	18.3	12,15	15	22.3,23.3	16	19	14,15	25,26	
AKR/J	JL STOCK#648	17	20.3	17	15	20.3	17	16	11	26	
ALR/ILT	JL STOCK# 3070	22	19.3	12	15	20.3	17.3	21	15	27	
BTBR T-> TF/J	JL STOCK# 2282	19	19.3	12	15	23.3	17.3	16	17	28	
C58/J	JL STOCK#669	16	20.3	15	15	22.3	14.3	18	16	25	
CE/J	JL STOCK#657	21	16.3	16	16	20.3	17	15	13	24	
CZECH/EJ	JL STOCK# 2799	18	18.3	18	7.1	14	17	17.1	15	25.3	
FVB/NJ	JL STOCK# 1800	17	19.3	12	15	20.3	14.3	20	14	26	
JF1/MS	JL STOCK# 3720	23	17	OUT OF RANGE	7.1	17	18	21.1	18	32	
MA/MyJ	JL STOCK#677	17	19.3,20.3	14	15	24.3	15.3	17	14	27	
MOLC/RK	JL STOCK# 731	16	17	13	7.1	17	18	24.1	18	32	
MOLF/EJ	JL STOCK# 550	15	17	17	7.1	17	18	24.1	14	32	
MRL/MPJ	JL STOCK#486	17	20.3	12	15	25.3	17	16	15	26	

FIG. 9 (Cont.)

MOUSE DNA SAMPLE	ORIGIN	MOUSE STR MARKERS									
		18-3	4-2	6-7	9-2	15-3	6-4	12-1	5-5	X-1	
MSM/MS	JL STOCK# 3719	18	18	26	7.1	17	20	18.1	21	37	
NON/SHILT	JL STOCK# 1290	17	20.3	15	15	22.3	14.3	16.1	13	25	
NOR/LTJ	JL STOCK# 2050	18	18.3	14	15	20.3	15.3	17	15	27	
NZW/LACJ	JL STOCK# 1058	17	17.3	15	15	22.3	17	18	11	25	
PL/J	JL STOCK# 680	18	19.3	12	15	22.3	17	16	14	26	
RIIS/J	JL STOCK# 683	18	19.3	14	15	25.3	16	17	15	25	
SB/LEJ	JL STOCK# 269	16	18.3	13	16	20.3	17	17	17	26	
SEA/GNJ	JL STOCK# 644	18	21.3	12	15	23.3	17	17	15,16	28	
SJL/J	JL STOCK# 686	17	19.3	12	15	22.3	14	17	14	26	
SPRET/EIJ	JL STOCK# 1146	OUT OF RANGE	OUT OF RANGE	17.3	15.1	OUT OF RANGE	11.2	17.1	17	17.1	
NIH3T3	ATCC#CRL-1658	17,19	19.3	12	15,16	20.3	14.3	20	14,15	25	
L929	ATCC#CCL-1	16	20.3	12	15	24.3,25.3,26.3	16	16	14	26,27	
MC3T3-E1 SUBCL 4	ATCC#CRL-2593	15	20.3	17	17,18	22.3	17	17	17	28	
W-20-17	ATCC#CRL-2623	16,18	20.3,23.3	12,17	18	22.3	17	16,17	12,17	27	
HK-PEG-1	ATCC#CCL-189	18,19	21.3	12	15	22.3,23.3	17,18	16	13,14	25	
P3X63A8.653	ATCC#CRL-1580	18,19	21.3	12	15,16	22.3,23.3	17,18	16	13,14	25	
RAW 264.7	ATCC#TIB-71	18	22.3	12	15	22.3	17	16	14	24	
M. DUNNI	ATCC#CRL-2017	15	17.1	20.1,23.1	11.1	12	NO AMPLIF	17,19,21	13,16	OUT OF RANGE	

FIG. 9 (Cont.)

Sequence ID No.	STR Marker	NCBI GenBank/ Accession#	Primers (5'-3')	Primer $\mu$ M	Repeat Motif
97	Mo11-1	AC_000033.1 (62179023-62179299)	F4: [6FM]-CCAGCAGGCACTCTAGTTA R4: GTTCTTACAGCTTCAGGACATGGAA	0.200	[ATAC] <sub>n</sub> [R] <sub>n</sub> [G] <sub>n</sub> [AG] <sub>n</sub>
98	Mo19-2	AC_000041.1 (3931717-3932251)	F4: [6FM]-AGCTAGCACTGTCCTTGT R7: GTTCTTACTCAGCACCCTCCATCCTG	0.200	[CTAT] <sub>n</sub>
99	Mo1-2	AC_000023.1 (4308426-4308549)	F3: [VIC]- TCTTAAAAATCAAAACAGGCAAA R: GTTCTTGGGAGGTTGGGGTGATAA	0.200	[TAGA] <sub>n</sub> [TGAT] <sub>n</sub> [AGAT] <sub>n</sub>
100	Mo7-1	NW_004058052.1 (409093-409299)	F: [VIC]-ACACCAGTTGGGGACTAGGA R: GTTCTTAACTGGTGCCTGGGCTGAAC	0.200	[ATC] <sub>n</sub> [TTAT] <sub>n</sub> [NL] <sub>14</sub> [TC] <sub>n</sub> [N2] <sub>6</sub> [ATT] <sub>n</sub>
101	Mo1-1	NC_000067.6 (169358615-169358969)	F: [VIC]-CCCTTCACTCCTTCATTCA R: GTTCTTGGACCTAAGGACCTGGACA	0.200	[ATAC] <sub>n</sub>
102	Mo3-2	AC_000025.1 (3607722-3608165)	F3: [VIC]-TGAGTACCATGTGGGTACA R3: GTTCTTCACACACACACAAAGATGGA	0.200	[TATC] <sub>n</sub>
103	Mo8-1	AC_000030.1 (4494304-4494841)	F4: [VIC]-AGTAAATCCTGGTCTGGGC R4: GTTCTTGGCTCAGTATGAGCTATTGGA	0.200	[TATG] <sub>n</sub>
104	Mo2-1	AC_000024.1 (125349559-125349718)	F2: [NED]-ACAACCTCAAACTTGGCTGGAA R3: GTTCTTCCCTGCTAGCTTGCATTCC	0.100	[ATAC] <sub>n</sub>

FIG. 10

105	Mo11-2	NC_000077.6 (3318994-3319352)	F2: [NED]-AAGGCAGGGGAATTCACAGT R4: GTTCTTCTCACCATTGCAGTCTGA	0.200	[ATCT] <sub>n</sub>
106	Mo17-2	AC_000039.1 (5594131-5594332)	F: [PET]-CTCTTCCATCGTCCATC R: GTTCTTATGGCTCTGCCACAATCT	0.200	[CTAT] <sub>n</sub>
107	Mo13-1	AC_000035.1 (3646204-3646733)	F4: [PET]-AACTCCCACTGCTTCTGGT R6: GTTCTTGCCAAGTAAACCCTTCTCCC	0.250	[TCTA] <sub>n</sub>

FIG. 10(Cont.)

(SEQ ID NO. 60)

STR Marker	Primers (5'-3')	Amplicon Size (bp)	T <sub>a</sub> (C°)
Mo11-1	FSeq1: CACCCATGAAAGAGCCCAAGT*; RSeq1: TTCAGGACCATGGAAGCTC* (SEQ ID NO. 61)	342-367	60
Mo19-2	FSeq1: TCCTCAAGGCTAGCACTGTTCT* (SEQ ID NO. 62) RSeq1: AAATCCCAGCACTCAGGAGA* (SEQ ID NO. 63)	276-300	60
Mo1-2	F: ACTCCTCCCAGATTCTTCC* ; (SEQ ID NO. 64) RSeq2: TGCCCAATGTATGGTTTCC* (SEQ ID NO. 65)	199-236	60
Mo7-1	F: ACACCAGTTGGGACTAGGA ; (SEQ ID NO. 66) R3: CTGCTCTCCAGAGGTCCTG* (SEQ ID NO. 67)	349-405	60
Mo1-1	F: CCCTTCACTCCTTCATTCCA ; (SEQ ID NO. 68) R: TGAGCCTAAGGACCTGGACA (SEQ ID NO. 69)	325-366	59
Mo3-2	F: CCATGTGGGTACATIGTAAAGC* ; (SEQ ID NO. 70) R: CATGTGATAGAGGATGGATAGTA* (SEQ ID NO. 71)	277-347	59
Mo8-1	F: TGGCAGAGTGCATACCTAGC* ; (SEQ ID NO. 72) R: CTAGAACTGGGCTGGTGGT* (SEQ ID NO. 73)	271-325	59
Mo2-1	FSeq1: CCTAGAGTCTTGTGGCTGGA* ; (SEQ ID NO. 74) RSeq1: CACCTAGTCTCCGATTGA* (SEQ ID NO. 75)	373-408	60
Mo11-2	F: TCCAGGGGAATCTTTGACAC* (SEQ ID NO. 76) R: GAGGGCTGTAAGTTGTGC* (SEQ ID NO. 77)	203-260	59
Mo17-2	F: CTCTTCTCCATCCGTCCTC* ; (SEQ ID NO. 78) R: ATGGCTCTTGCCACAAAATCT (SEQ ID NO. 79)	183-227	59
Mo13-1	F: TCGTTACCAATGCCCTCTTCC* ; (SEQ ID NO. 80) R3: CGCCACAGCCTCATCTACTC* (SEQ ID NO. 81)	309-345	60

FIG. 11

FRAGMENT LENGTH IS ABOVE THE REPEAT NUMBER WHICH IS IN PARENTHESIS		ALLELE DISTRIBUTION AND CORRELATING FRAGMENT LENGTH (BP)				
STR MARKER	REPEAT MOTIF	266	268	269	272	273
MO11-1	[ATAC] <sub>N</sub> [R] <sub>1</sub> [G] <sub>N</sub> [AG] <sub>N</sub>	(17)	(17.2)	(17.3)	(18.2)	(18.3)
MO19-2	[CTAT] <sub>N</sub>	533	537	540	545	549
MO1-2	[TAGA] <sub>N</sub> [TGAT] <sub>N</sub> [AGAT] <sub>N</sub>	(10)	(11)	(12)	(13)	(14)
MO7-1	[ATCT] <sub>N</sub> [TTIAT] <sub>N</sub> [N1] <sub>4</sub> [TC] <sub>N</sub> [N2] <sub>6</sub> [ATTT] <sub>N</sub>	104	108	112	116	120
		(12)	(13)	(14)	(15)	(16)
		183	195	199	203	205
		(19.2)	(22.2)	(23.2)	(24.2)	(25)
MO1-1	[ATAC] <sub>N</sub>	330	334	338	342	346
		(10)	(11)	(12)	(13)	(14)
MO3-2	[TATC] <sub>N</sub>	431	435	439	443	447
		(9)	(10)	(11)	(12)	(13)
MO8-1	[TATG] <sub>N</sub>	503	527	530	531	535
		(6)	(12)	(12.3)	(13)	(14)
MO2-1	[ATAC] <sub>N</sub>	124	128	132	136	140
		(8)	(9)	(10)	(11)	(12)
MO11-2	[ATCT] <sub>N</sub>	407	411	413	415	417
		(12)	(13)	(13.2)	(14)	(14.2)
MO17-2	[CTAT] <sub>N</sub>	188	192	196	200	202
		(11)	(12)	(13)	(14)	(14.2)
MO13-1	[TCTA] <sub>N</sub>	511	519	523	527	528
		(11)	(13)	(14)	(15)	(15.1)

FIG. 12



274	275	276	277	278	279	280	281	282	283	285	288	289	293
(19)	(19.1)	(19.2)	(19.3)	(20)	(20.1)	(20.2)	(20.3)	(21)	(21.1)	(21.3)	(22.2)	(22.3)	(23.3)
552	553	557											
(14.3)	(15)	(16)											
122	124	128	132	136	140	145							
(16.2)	(17)	(18)	(19)	(20)	(21)	(22)							
207	209	211	213	215	217	219	221	225	229	232	233	237	239
(25.2)	(26)	(26.2)	(27)	(27.2)	(28)	(28.2)	(29)	(30)	(31)	(31.3)	(32)	(33)	(33.2)
350	354	358	362	366									
(15)	(16)	(17)	(18)	(19)									
448	451	455	459	463	467	471	495						
(13.1)	(14)	(15)	(16)	(17)	(18)	(19)	(25)						
539	543	547	551	554	555								
(15)	(16)	(17)	(18)	(18.3)	(19)								
144	148	152	156	160									
(13)	(14)	(15)	(16)	(17)									
419	422	423	426	427	429	430	431	432	435	438	439	465	
(15)	(15.3)	(16)	(16.3)	(17)	(17.2)	(17.3)	(18)	(18.1)	(19)	(19.3)	(20)	(26.2)	
204	205	208	212	216	224	232							
(15)	(15.1)	(16)	(17)	(18)	(20)	(22)							
529	531	532	533	535	536	537	539	540	541	543	544	546	548
(15.2)	(16)	(16.1)	(16.2)	(17)	(17.1)	(17.2)	(18)	(18.1)	(18.2)	(19)	(19.1)	(19.3)	(20.1)

FIG. 12 (Cont.)



NEW PRIMERS AND ALLELE DISTRIBUTION OF CHANGES TO THE ORIGINAL 9STR MARKERS (TO ACCOMMODATE THEM INTO THE NEW 21PLEX)												
ALLELE DISTRIBUTION												
15-3	AVG BP	178	186	190	193	194	198	201	202	205	206	
	REPEAT	12	14	15	15.3	16	17	17.3	18	18.3	19	
F: [NED]-TCTGGCGTGTCTGCATAA (SEQ ID NO. 87)												
R-EPIC: GTTTCCTTAGGCTATCTAAACTTGACATGCT (SEQ ID NO. 88)												
ALLELE DISTRIBUTION												
X-1	AVG BP	386	392	393	396	400	404	408	412	416	420	
	REPEAT	15.2	17	17.1	18	19	20	21	22	23	24	
F-B: [PET]-AACAAAAATGTCCTCAATGC (SEQ ID NO. 93)												
R3PIG: GTTTCCTTAGGCTATCTAAACTTGACATGCT (SEQ ID NO. 94)												
ALLELE DISTRIBUTION												
12-1	AVG BP	264	268	269	272	273	276	277	280	281	284	
	REPEAT	15	16	16.1	17	17.1	18	18.1	19	19.1	20	
F2: [PET]-TTTCAAAAATGTCATTGAACACA (SEQ ID NO. 89)												
R6PIG: GTTTCCTTAGGCTATCTAAACTTGACATGCT (SEQ ID NO. 90)												
D8S1106 (HUMAN STR)												
F2: TCTGCTGATTGTGCCCATTTG (SEQ ID NO. 108)												
R3PIG: GTTTCCTTAGGCTATCTAAACTTGACATGCT (SEQ ID NO. 109)												

FIG. 14

209	213.42	217	221	225	229	233	237	241	243	245	258
<b>19.3</b>	<b>20.3</b>	<b>21.3</b>	<b>22.3</b>	<b>23.3</b>	<b>24.3</b>	<b>25.3</b>	<b>26.3</b>	<b>27.3</b>	<b>28.1</b>	<b>28.3</b>	<b>31.3</b>
424	427	428	431	435	439	443	451	471			
<b>25</b>	<b>25.3</b>	<b>26</b>	<b>27</b>	<b>28</b>	<b>29</b>	<b>30</b>	<b>32</b>	<b>37</b>			
288	289	292	296	300	301						
<b>21</b>	<b>21.1</b>	<b>22</b>	<b>23</b>	<b>24</b>	<b>24.1</b>						

FIG. 14 (Cont.)

**MOUSE CELL LINE AUTHENTICATION****CROSS-REFERENCE TO RELATED APPLICATIONS**

The present application claims priority to provisional application Ser. No. 61/692,761, filed on or about Aug. 24, 2012, entitled "Mouse Cell Line Authentication" naming the same inventors as in the present application. The present application also claims priority to nonprovisional application Ser. No. 13/935,285 filed on or about Jul. 3, 2013, entitled "Mouse Cell Line Authentication," naming the same inventors as the present application. The contents of both this provisional and nonprovisional application are incorporated by reference herein, the same as if fully set forth.

**STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH/DEVELOPMENT**

The subject matter of this patent application was invented under the support of at least one United States Government contract. Accordingly, the United States Government may manufacture and use the invention for governmental purposes without the payment of any royalties.

**SEQUENCE LISTING**

The present application has a sequence listing set forth in a computer-readable file submitted herewith that is titled as follows: "NIST10005016CIP\_ST25final\_May 2020". Said computer readable file has a size of twenty-eight (28) kilobytes and is hereby incorporated by reference in the present application, the same as if fully set forth.

**BACKGROUND OF THE INVENTION****Field of Invention**

The present disclosure relates to cell lines and mouse strains, more particularly, to mouse cell line authentication.

**Description of Related Art**

Cell line authentication is becoming increasingly important. For example, cell line authentication is now required by some journals prior to publication. In some cases, cell line authentication may be mandatory before receiving funding from some granting agencies.

The Federal Drug Administration has also instituted a requirement for the authentication of cell lines used to produce pharmaceuticals. Methods are currently in place for authenticating human cell lines using multiplex PCR assays that target short tandem repeat (STR) markers in the human genome and are capable of generating a unique individual genotypic profile. Cell repositories may now genotype their human cell lines using at least eight human STR markers including a marker for amelogenin, the sex identification locus.

Large databases of STR profiles may be used to confirm genotypes of human cell lines. These databases may also be used to provide a record of previously misidentified or cross-contaminated human cell lines. The American Type Culture Collection (ATCC) website maintains an updated list of misidentified human cell lines. Although there are successful methods in place for human cell line authentication, methods for nonhuman cell lines are not well established.

Mouse cell lines are commonly used to study human genes and disease. Mouse cells may also be used in the

biomanufacturing of recombinant proteins and may also function as feeder cells for embryonic stem cells.

In the prior art, there are a number of techniques used to identify mouse cell lines or mouse strains, including but not limited to microsatellite markers (simple sequence length polymorphism (SSLP) or STR markers), single nucleotide polymorphisms (SNPs), and species-specific primers. One of the most commonly used laboratory strains is the *Mus musculus domesticus* strain. This strain may be difficult to genotype due to many shared alleles as a result of extensive breeding.

There have been published reports of multiplexing mouse SSLPs post-PCR by pooling the amplified products to distinguish between different strains of inbred mice. However, most of the microsatellite markers that have been used for these purposes are dinucleotide in nature, mainly CA repeats, which may result in noisy stutter and have relatively high mutation frequencies when compared to tetranucleotide repeats. The reduced stutter associated with tetranucleotide repeats may allow for easier interpretation of single and mixed profiles.

There is a need for mouse cell line authentication based on tetranucleotide repeats, as opposed to microsatellite markers that are dinucleotide in nature.

Current methods may lack the resolution to differentiate between individual mice of the same subspecies. SNPs may be well-conserved between inbred mice of the same strain. Thus, it may be difficult to differentiate between interstrain mice using this method. Even an extensive array containing over 600,000 mouse SNPs may still be unable to identify individual mice within the same subspecies.

Species-specific primers may be used to determine the origin of species for cell lines. However, species-specific primers may lack specificity to identify at the individual level.

For human cell lines, practices have been adopted that are currently used to identify such cells. These practices are based on the detection of short tandem repeat (STR) markers. However, for the large volume of research that is performed using nonhuman cell lines (mouse, hamster, monkey, etc.), there are no STR marker assays that are able to uniquely and unequivocally identify a particular cell line.

There is further a need for a mouse cell line authentication assay that uniquely identifies cell lines at the individual level.

**BRIEF SUMMARY OF DISCLOSURE**

The present disclosure addresses the needs described above by providing a PCR assay for mouse cell line authentication. The present assay may be used to authenticate mouse cell lines resulting in unique profiles for individual mouse samples based on tetranucleotide repeats that are stable with high passage number in the two different cell lines tested.

In accordance with one embodiment of the present disclosure, a method is provided for determining the alleles present in a DNA sample. The method comprises obtaining a DNA sample to be analyzed; and selecting a set of STR marker loci of the DNA sample to be analyzed that can be amplified together in a multiplex amplification reaction, wherein the set of STR marker loci are selected from the group consisting of: 18-3, 4-2, 6-7, 15-3, 6-4, 12-1, 5-5, X-1, Mo11-1, Mo19-2, Mo1-2, Mo7-1, Mo1-1, Mo3-2, Mo8-1, Mo2-1, Mo11-2, Mo17-2, Mo13-1, D8S1106 and D4S2408.

The method further comprises providing a set of oligonucleotide primer pairs, wherein each oligonucleotide

primer pair in the set flanks a single locus in the set of STR marker loci, and wherein each primer pair is capable of amplification of a single locus from the set of STR marker loci in a multiplex amplification reaction. The method also comprises co-amplifying the set of STR marker loci in a multiplex amplification reaction, wherein the product of the multiplex amplification reaction comprises a mixture of amplified alleles from each of the co-amplified loci in the set of STR marker loci. Finally, the method comprises evaluating the products of the co-amplification reaction to determine the alleles present at each of the loci analyzed in the set of STR marker loci within the DNA sample.

In accordance with another embodiment of the present disclosure, a kit is provided for determining the alleles present in a DNA sample to be analyzed. The kit comprises oligonucleotide primer pairs for co-amplifying a set of STR marker loci of at least one DNA sample to be analyzed. The set of loci comprises one or more of STR loci selected from the group consisting of STR marker loci 18-3, 4-2, 6-7, 15-3, 6-4, 12-1, 5-5, X-1, Mo11-1, Mo19-2, Mo1-2, Mo7-1, Mo1-1, Mo3-2, Mo8-1, Mo2-1, Mo11-2, Mo17-2, Mo13-1, D8S1106 and D4S2408.

These, as well as other objects, features and benefits will now become clear from a review of the following detailed description of illustrative embodiments and the accompanying drawings.

#### BRIEF DESCRIPTION OF DRAWINGS

FIG. 1 is a table illustrating primers for STR amplification and their corresponding chromosomal locations in accordance with the present disclosure.

FIG. 2 is a table illustrating sequencing primers and their respective amplicon size range as well as annealing temperatures in accordance with the present disclosure.

FIG. 3 is a table defining STR fragment length and corresponding repeat number in accordance with the present disclosure.

FIG. 4 is a table illustrating the complete genetic profiles of six mouse cell lines in accordance with the present disclosure.

FIG. 5 is a genetic profile of an NIH3T3 cell line using the multiplex assay of the present disclosure.

FIG. 6 illustrates the results obtained when a human contaminant is detected in an NIH3T3 STR profile in accordance with one embodiment of the present disclosure.

FIG. 7 is a genetic profile of a RAW 264.7 cell line using the multiplex assay of the present disclosure.

FIG. 8 is a mixture of NIH3T3 and RAW 264.7 cell lines detected using the multiplex assay of the present disclosure.

FIG. 9 is a table containing the STR profiles showing the allele ranges for the markers from mouse cell lines, mouse strains, and wild mice.

FIG. 10 is another table illustrating primers for STR amplification and their corresponding chromosomal locations in accordance with one embodiment of the present disclosure.

FIG. 11 is another table illustrating sequencing primers and their respective amplicon size range as well as annealing temperatures in accordance with one embodiment the present disclosure.

FIG. 12 is a table defining STR fragment length and corresponding repeat number in accordance with one embodiment of the present disclosure.

FIG. 13 includes updated primer sequences for various STR markers in accordance with one embodiment of the present disclosure.

FIG. 14 illustrates four updated primers and allele distribution of changes to the various STR markers (to accommodate them into the new 21-plex assay).

#### DEFINITIONS

**Allele:** The term “allele” as used herein refers to a genetic variation associated with a gene or a segment of DNA, i.e., one of two or more alternate forms of a DNA sequence occupying the same locus.

**Amplicon/amplification product/amplified sequence:** The terms “amplicon,” “amplification product” and “amplified sequence” are used interchangeably herein and refer to a broad range of techniques for increasing polynucleotide sequences, either linearly or exponentially and can be the product of an amplification reaction. An amplicon can be double-stranded or single-stranded, and can include the separated component strands obtained by denaturing a double-stranded amplification product. In certain embodiments, the amplicon of one amplification cycle can serve as a template in a subsequent amplification cycle. Exemplary amplification techniques include, but are not limited to, PCR or any other method employing a primer extension step. Other nonlimiting examples of amplification include, but are not limited to, ligase detection reaction (LDR) and ligase chain reaction (LCR). Amplification methods can comprise thermal-cycling or can be performed isothermally. In various embodiments, the term “amplification product” and “amplified sequence” includes products from any number of cycles of amplification reactions.

**Amplify:** As used herein, “amplify” refers to the process of enzymatically increasing the amount of a specific nucleotide sequence. This amplification is not limited to but is generally accomplished by PCR. As used herein, “denaturation” refers to the separation of two complementary nucleotide strands from an annealed state. Denaturation can be induced by a number of factors, such as, for example, ionic strength of the buffer, temperature, or chemicals that disrupt base pairing interactions. As used herein, “annealing” refers to the specific interaction between strands of nucleotides wherein the strands bind to one another substantially based on complementarity between the strands as determined by Watson-Crick base pairing. It is not necessary that complementarity be 100% for annealing to occur. As used herein, “extension” refers to the amplification cycle after the primer oligonucleotide and target nucleic acid have annealed to one another, wherein the polymerase enzyme catalyzes primer extension, thereby enabling amplification, using the target nucleic acid as a replication template.

**Detecting:** The terms “detecting” and “detection” are used in a broad sense herein and encompass any technique by which one can determine the presence of or identify a nucleic acid sequence. In some embodiments, detecting comprises quantitating a detectable signal from the nucleic acid, including without limitation, a real-time detection method, such as quantitative PCR (“Q-PCR”). In some embodiments, detecting comprises determining the sequence of a sequencing product or a family of sequencing products generated using an amplification product as the template; in some embodiments, such detecting comprises obtaining the sequence of a family of sequencing products. In other embodiments detecting can be achieved through measuring the size of a nucleic acid amplification product.

**Locus-specific allelic DNA size marker:** The term “locus-specific allelic DNA size marker” as used herein refers to a nucleic acid size standard for one or more alleles for a particular STR locus or marker. Those of skill in the art may

variably refer to this as an “allelic ladder.” The allelic ladder serves as a reference standard and nucleic acid size marker for the amplified alleles from the locus. In some embodiments, the allelic ladder can comprise size standards for the alleles of different STRs. In some embodiments, the allelic ladder can be made of DNA. In some embodiments, the allelic ladder can be made of non-naturally occurring nucleic acid analogs. The different individual size standards within an allelic ladder can, in some embodiments, be labeled with a detectable label, e.g., a fluorophore. In some embodiments, the allelic ladder components are labeled with the same fluorophore. In some embodiments, the allelic ladder components are labeled with different fluorophores. The size standards can be selected to work for a specific pair (or pairs) of oligonucleotide primers. For example, if a first set of primers for marker X with a tetranucleotide repeat produces a 150 base pair amplicon corresponding to allele 1, the corresponding allelic ladder component will serve as a size standard for the 150 base amplicons; while a second pair of primers for marker X produces a 154 base pair amplicon corresponding to allele 2, the corresponding allelic ladder component will serve as a size standard for the 154 base amplicons. Thus different size standards for different size amplicons of the same marker are contemplated. The size standard for a given amplicon derived from a given allele may have nucleic acid base sequence that is the same or different than the nucleic acid base sequence of the amplicon or allele from which the amplicon is derived. For allele analysis in electrophoresis systems the size standard can be selected so as to have the same electrophoretic mobility as the amplicon of interest. Alternatively, in some embodiments, the size standard can be selected so as to have different electrophoretic mobility than the amplicon of interest, given an understanding of the predictable nature of the difference; the identity of the amplicons could be determined. For allele analysis in mass spectroscopy systems the size standard (weight/charge ratio, not electrophoretic mobility) can be selected so as to have the same signal as the amplicon of interest. Alternatively, in some embodiments, the size standard (weight/charge ratio, not electrophoretic mobility) can be selected so as to have the different separation properties than the amplicon of interest, given an understanding of the predictable nature of the difference, the identity of the amplicons could be determined.

**Primer:** The term “primer” refers to a polynucleotide (oligonucleotide) and analogs thereof that are capable of selectively hybridizing to a target nucleic acid or “template”, a target region flanking sequence or to a corresponding primer-binding site of an amplification product; and allows the synthesis of a sequence complementary to the corresponding polynucleotide template, flanking sequence or amplification product from the primer’s 3’ end. Typically a primer can be between about 10 to 100 nucleotides in length and can provide a point of initiation for template-directed synthesis of a polynucleotide complementary to the template, which can take place in the presence of appropriate enzyme(s), cofactors, substrates such as nucleotides (dNTPs) and the like.

**Amplification Primer/Oligonucleotide primer:** As used herein, the terms “amplification primer” and “oligonucleotide primer” are used interchangeably and refer to an oligonucleotide, capable of annealing to an RNA or DNA region adjacent a target sequence, and serving as an initiation primer for DNA synthesis under suitable conditions well known in the art. Typically, a PCR reaction employs an “amplification primer pair” also referred to as an “oligonucleotide primer pair” including an “upstream” or “for-

ward” primer and a “downstream” or “reverse” primer, which delimit a region of the RNA or DNA to be amplified. A first primer and a second primer may be either a forward or reverse primer and are used interchangeably herein and are not to be limiting.

**STR Marker/STR Locus:** As used herein the terms “STR marker” and “STR locus”, and their plural forms, are used to describe one of a set of marker loci suitable for use in genotyping using the method of the present invention. In one embodiment:

“18-3” refers to the STR marker on mouse chromosome 18, corresponding to base pairs 60271556-60271705 (NCBI 38.1 mouse build, corresponding to GenBank Accession # NT\_039674.8) (REPEAT MOTIF: [ATCT]<sub>n</sub>);

“4-2” refers to the STR marker on mouse chromosome 4, corresponding to base pairs 82068280-82068580 (NCBI 38.1 mouse build, corresponding to GenBank Accession # NT\_187032.1) (REPEAT MOTIF: [GATA]<sub>n</sub>[GATG]<sub>n</sub>[A-TAG]<sub>n</sub>);

“6-7” refers to the STR marker on mouse chromosome 6, corresponding to base pairs 51601265-51601685 (NCBI 38.1 mouse build, corresponding to GenBank Accession # NT\_039353.8) (REPEAT MOTIF: [CTAT]<sub>n</sub>);

“9-2” refers to the STR marker on mouse chromosome 9, corresponding to base pairs 74395400-74395000 (NCBI 38.1 mouse build, GenBank Accession # NT\_039474.8) (REPEAT MOTIF: [TAGA]<sub>n</sub>[AGAT]<sub>n</sub>);

“15-3” refers to the STR marker on mouse chromosome 15, corresponding to base pairs 4930200-4930500 (NCBI 38.1 mouse build, GenBank Accession # NT\_039617.8) (REPEAT MOTIF: [TAGA]<sub>n</sub>);

“6-4” refers to the STR marker on mouse chromosome 6, corresponding to base pairs 142021975-142022270 (NCBI 38.1 mouse build, GenBank Accession # NT\_039360.8) (REPEAT MOTIF: [ATAG]<sub>n</sub>[ATGA]<sub>n</sub>[TAGA]<sub>n</sub>);

“12-1” refers to the STR marker on mouse chromosome 12, corresponding to base pairs 38480950-38481170 (NCBI 38.1 mouse build, GenBank Accession # NT\_039548.8) (REPEAT MOTIF: [AGAT]<sub>n</sub>[GATA]<sub>n</sub>);

“5-5” refers to the STR marker on mouse chromosome 5, corresponding to base pairs 112641540-112641821 (NCBI 38.1 mouse build, GenBank Accession # NT\_109320.5) (REPEAT MOTIF: [TATC]<sub>n</sub>); and

“X-1” refers to the STR marker on mouse chromosome X, corresponding to base pairs 110959842-110960080 (NCBI 38.1 mouse build, GenBank Accession # NT\_039706.8) (REPEAT MOTIF: [ATAG]<sub>n</sub>[ATGA]<sub>n</sub>[TAGA]<sub>n</sub>).

In another embodiment, as used herein the terms “STR marker” and “STR locus”, and their plural forms, are used to describe one of a set of marker loci suitable for use in genotyping using the method of the present disclosure. In particular:

“Mo11-1” refers to the STR marker on mouse chromosome 11, corresponding to base pairs 62179023-62179299 (GRCh37.p13 Primary Assembly mouse build, corresponding to GenBank Accession # AC\_000033.1)(REPEAT MOTIF: [ATAC]<sub>n</sub>[R]<sub>n</sub>[G]<sub>n</sub>[AG]<sub>n</sub>), where R is any purine (A or G);

“Mo19-2” refers to the STR marker on mouse chromosome 19, corresponding to base pairs 3931717-3932251 (GRCh37.p13 Primary Assembly mouse build, corresponding to GenBank Accession # AC\_000041.1) (REPEAT MOTIF: [CTAT]<sub>n</sub>);

“Mo1-2” refers to the STR marker on mouse chromosome 1, corresponding to base pairs 4308426-4308549 (GRCh37.p13 Primary Assembly mouse build, correspond-





Two well-characterized human STR markers, D8S1106 and D4S2408, may be included in the multiplex kit (sometimes referred to hereinafter as an “assay”). These markers may be used to screen for contamination of mouse cell lines with human or African green monkey cell lines. Both human STR markers can be used to identify human and African green monkey cell lines (e.g., Vero cells).

Primer3 software, available online from the Massachusetts Institute of Technology, may be used to design PCR primers to flank the STR regions by inputting the downloaded mouse sequences from the NCBI BLAST program. The parameters for Primer3 were set to target primers with annealing temperatures of 60° C. AUTODIMER™ software was used to assess primer-dimer interactions and hairpin structures of possible primer combinations to be used in the multiplex. Forward primers were labeled with one of the following fluorescent dyes at the 5' end: 6FAM™ (blue), VIC™ (green), NED™ (yellow), or PET™ (red) (Applied Biosystems, Foster, Calif.). In some cases, an additional guanine base (G) or a “PIGtail” sequence (GTTTCTT) was added to the 5' end of the unlabeled reverse primers to promote complete adenylation.

Referring now to FIG. 1, illustrated is a table showing the primers used for STR amplification and their corresponding chromosomal locations. In FIG. 1, mouse chromosomal locations, or base pairs (bp), are based on NCBI BLAST 38.1 mouse build. Chromosomal locations for human STR markers D8S1106 and D4S2408 are based on the NCBI 37.3 human build. Primer concentrations in the rightmost column are final concentrations of forward and reverse primers in a 20 µL reaction volume. Primer concentrations were determined empirically based on peak height, DNA concentration and the number of cycles in the PCR program.

Based on FIG. 1, the exemplary amplification primers are set forth below. In order to specifically amplify the selected STR markers, the following oligonucleotide primer pairs were used, where “F” and “R” correspond to the forward and reverse primers respectively. The PCR is the product expected from amplification of the particular STR locus to which the oligonucleotide pair was directed.

18-3:	(SEQ ID NO: 10)	
F-TCTTTCTCCTTTTGTGTCATGC		
	(SEQ ID NO: 11)	
R-GTTTCTTGCTAAATAACTAAGCAAGTGAACAGA		
Primer	(SEQ ID NO: 1)	
4-2:	(SEQ ID NO: 12)	
F-AAGCTTCTCTGGCCATTTGA		
	(SEQ ID NO: 13)	
R-GTTCATAAACTTCAAGCAATGACA		
Primer	(SEQ ID NO: 2)	
6-7:	(SEQ ID NO: 14)	
F-AGTCCACCCAGTGCATTCTC		
	(SEQ ID NO: 15)	
R-GTTTCTTCATGTGGCTGGTATGCTGTT		
Primer	(SEQ ID NO: 3)	

-continued

9-2:	(SEQ ID NO: 16)	
F-GGATTGCCAAGAATTTGAGG		
	(SEQ ID NO: 17)	
R-GTTTCTTTCTCCTGAGTTGTGGACAGGGTTA		
Primer	(SEQ ID NO: 4)	
15-3:	(SEQ ID NO: 18)	
F-TCTGGGCGTGTCTGTCATAA		
	(SEQ ID NO: 19)	
R-GTTTCTTTTCTCAGGGAGGAGTGTGCT		
Primer	(SEQ ID NO: 5)	
6-4:	(SEQ ID NO: 20)	
F-TTTGCAACAGCTCAGTTTCC		
	(SEQ ID NO: 21)	
R-GTTTCTTAATCGCTGGCAGATCTTAGG		
Primer	(SEQ ID NO: 6)	
12-1:	(SEQ ID NO: 22)	
F-CAAATTGTCATTGAACACATGTAA		
	(SEQ ID NO: 23)	
R-GTTTCTTTCAATGGTCAAGAAATACTGAAGTACAA		
Primer	(SEQ ID NO: 7)	
5-5:	(SEQ ID NO: 24)	
F-CGTTTTACCTGGCTGACACA		
	(SEQ ID NO: 25)	
R-GTTTCTTTGGTTTAAAACTCAATACCAACAA		
Primer	(SEQ ID NO: 8)	
X-1:	(SEQ ID NO: 26)	
F-GGATGGATGGATGGATGAAA		
	(SEQ ID NO: 27)	
R-GTTTCTTAAGGTATATATCAAGATGGCATTATCA		
Primer	(SEQ ID NO: 9)	
D8S1106:	(SEQ ID NO: 30)	
F-GTTTACCCCTGCATCATGG		
	(SEQ ID NO: 31)	
R-GTTTCTTTCAGAATTGCTCATAGTGCAAGA		
Primer	(SEQ ID NO: 28)	
D4S2408:	(SEQ ID NO: 32)	
F-TCATTTCCATAGGGTAAGTGAAAA		
	(SEQ ID NO: 33)	
R-GTTTCTTGCCATGGGGATAAAATCAGA		
Primer	(SEQ ID NO: 29)	

Genomic mouse DNA samples were obtained from Jackson Laboratories (Bar Harbor, Me., USA). These samples

represent 48 common inbred strains used in the scientific community. The DNA included thirty-seven inbred mice DNA samples, one recombinant inbred mouse sample, and ten wild-derived mice DNA samples. DNA from 15 wild-caught mice (courtesy of Dr. Michael Nachman from the University of Arizona, USA) collected in Tucson, Ariz. was used for heterozygosity studies.

Genomic DNA from mouse (male and female CD1/ICR), hamster (Syrian golden hamster, Chinese hamster), rat (Fischer, Wistar, Sprague Dawley), gerbil, pig, baboon, rhesus, and cynomolgus monkey were obtained from Zyagen (San Diego, Calif.). TN1 cells (stably transfected green fluorescent protein (GFP) expressing cell line derived from the parent NIH3T3 line) were originally obtained from the American Type Culture Collection, Manassas, Va., in 2003. The following cell lines were obtained from The American Type Culture Collection (ATCC, Manassas, Va.): NIH3T3 (CRL-1658), L-929 (CCL-1), MC3T3-E1 subclone 4 (CRL-2593), RAW 264.7 (TIB-71), *M. dummi* (CRL-2017), P3X63Ag8.653 (CRL-1580), HK-PEG-1 (CCL-189), Vero (CCL-81), HeLa (CCL-2), and CHO-K1 (CCL-61).

DNA was quantified using the SYNERGY™ Mx plate reader and TAKE3™ plate (BioTek, Winooski, Vt., USA) at an absorbance of 260 nm. To study STR marker stability as passage number increased, duplicate 25 cm<sup>2</sup> tissue culture flasks of L929 cells were carried independently, and one million cells were harvested at passage numbers 2, 4, 9, 14, 19, 22, 26, 29, 31, 37, 41, and 44. Duplicate 25 cm<sup>2</sup> flasks were also carried for NIH3T3 cells which were carried independently, and one million cells were harvested from passage numbers 5, 7, 10, 15, 20, 23, 26, 32, 35, 40, 43, and 45.

PCR amplification was performed on a VERITI™ thermal cycler (Applied Biosystems). The reaction mixture of 20 μL final volume contained 1 ng of mouse DNA (or 5 ng to 10 ng of non-mouse DNA for specificity studies), 1× GENEAMP® PCR Gold buffer (Applied Biosystems), 2 mM MgCl<sub>2</sub> (Applied Biosystems), 250 μM dNTPs (USB Corporation, Cleveland, Ohio, USA), forward labeled and reverse primers (as shown in FIG. 1 hereinabove), 1U AMPLITAQ GOLD™ DNA Polymerase (Applied Biosystems), and 0.16 mg/mL non-acetylated BSA (Invitrogen). PCR conditions for the multiplex assay are as follows: denaturation for 11 min at 95° C., amplification for 30 cycles of 45 s at 94° C., 2 min at 59° C., and 1 min at 72° C., followed by an extension for 60 min at 60° C., and a final soak at 25° C.

Initial unlabeled primers and their respective PCR products were screened by using gel electrophoresis. PCR products (4 μL) were added to the Lonza 5X loading dye (1 μL), loaded onto a 2.2% agarose Flash Gel (Lonza) and run at 275 V for 5 min. Forward primers generating clean PCR products were ordered with a fluorescent dye at the 5' end and were tested in multiplex reactions with mouse DNA from Jackson Laboratories, Zyagen, and mouse cell lines. Multiplex reactions were then optimized by varying primer combinations, primer concentrations, DNA concentration, and PCR cycle number. To analyze monoplex and multiplex PCR products, samples were prepared by adding 1 μL of amplified product and 0.3 μL of GENESCAN™ 500 LIZ internal size standard (Applied Biosystems) to 8.7 μL of HI-DI™ formamide (Applied Biosystems) for separation on the 16-capillary ABI 3130xl Genetic Analyzer (Applied Biosystems). A five dye matrix was established under the G5 filter with dyes 6FAM, VIC, NED, PET, and LIZ. POP-4™ (Applied Biosystems) was used on a 36 cm capillary array (Applied Biosystems) with 1× ACE™ buffer (Amresco,

Solon, Ohio, USA). Samples were injected electrokinetically for 10 s at 3 kV. The STR alleles were separated at 15 kV at a run temperature of 60° C. Data from the 3130xl genetic analyzer was analyzed using the GENEMAPPER™ ID-X v1.1 Software (Applied Biosystems). Bins and panels were created in GENEMAPPER™ ID-X based on fragment length data generated from the fifty-seven mouse profiles using fixed bin allele sizes to determine allele calls. The allele distribution range for the human STR markers (D8S1106 and D4S2408) was previously described and adjustments were made to the size range to take into account the "PIGtail" sequence that was added to the reverse primers. Calibration of repeat number to allele fragment length was determined by DNA sequencing.

It should also be noted that while FIG. 1 refers to nine STR markers used for mouse cell line authentication, FIG. 10 herein below provides eleven (11) additional such STR markers.

Multiplex primers were used for sequencing STR markers, except for three loci (18-3, 9-2, and 12-1) where sequencing primers were used. Referring now to FIG. 2, illustrated are the forward and reverse primers used to sequence each of the nine STR markers. Also shown in FIG. 2 are the corresponding annealing temperatures and amplicon sizes for these markers.

At least four homozygous samples were sequenced for each STR locus to determine the corresponding number of repeats for each allele. The targeted repeat regions were amplified using 0.15 μM unlabeled forward and reverse primers using the PCR reaction specified herein in connection with PCR amplification with the following thermal cycling program: denaturation for 10 min at 95° C., amplification for 35 cycles of 1 min at 94° C., 1 min at 52-60° C. (annealing temperature specific to individual primers), and 1 min at 72° C., followed by an extension for 45 min at 60° C., and a final soak at 25° C. Samples were treated with 2 μL of EXOSAP-IT® PCR product cleanup (USB Corporation) per 5 μL of PCR product. This product cleanup was used to remove unincorporated primers and deoxyribonucleotide triphosphates (dNTPs) by incubating samples for 90 min at 37° C. followed by 20 min at 80° C. to inactivate the enzymes. Samples were then sent to Eurofins MWG Operon for sequencing using BIGDYE® Terminator v3.1 (Applied Biosystems). Resulting profiles were received after data analysis was performed by Eurofins MWG Operon.

Mixture samples containing genomic DNA extracted from NIH3T3, RAW264.7, and HeLa cells were analyzed to assess the capability of the multiplex assay to detect low levels of contamination in NIH3T3 cells. DNA from NIH3T3 and RAW264.7 cells were added to individual reactions with a final concentration of 1 nanogram (ng) of total DNA in the following ratios 1:1, 2:1, 3:1, 5:1, 7:1, 9:1, and 10:1. Reciprocal reactions were also prepared using DNA from RAW264.7 and NIH3T3 cells. The same procedure was repeated using DNA from NIH3T3 and HeLa cells, followed by reciprocal reactions with DNA from HeLa and NIH3T3 cells. PCR amplification and PCR product analysis are described above.

The heterozygosity (H) values were calculated by dividing the number of heterozygotes at a locus into the total number of individuals. The probability of identity (PI) was calculated by the summation of the square of the genotype frequencies. The probability of a random match (PM) for a full profile was calculated by multiplying the inverse of each genotype frequency for each marker. The coefficient of inbreeding (F), specifically the fixation in a subpopulation compared to the total population (FST) was determined by

subtracting the average heterozygosity of the two subpopulations (wild-caught mice and inbred mice samples) from the total heterozygosity, divided by the total heterozygosity.

The mouse primers targeting tetranucleotide repeat markers in the multiplex PCR assay were designed based on the annotated mouse genome from NCBI build 38.1 of *Mus musculus* origin. Fifty-seven genomic mouse DNA samples were tested using the multiplex assay and the designated allele range was determined for each marker, and fragment lengths were correlated to actual number of repeats using sequence analysis.

Referring now to FIG. 3, illustrated is a table defining STR fragment length and corresponding repeat number in accordance with the present disclosure. In FIG. 4, fragment length in base pairs corresponds to apparent size based on LIZ GENESCAN™ 500 size standard. The corresponding number of repeats are each shown just below the fragment length. The corresponding number of repeats was determined by the analysis of 57 mouse DNA samples. The correlation of the allele size and number of repeats was determined based on sequencing data.

The mouse samples were selected to represent the genetic diversity of the mouse family tree. To determine the specificity of the multiplex assay, DNA was tested from several different species and subspecies of mice, near neighbors, and non-mouse samples. A panel of 57 mouse genomic DNA samples representing species from *M. musculus musculus*, *M. musculus domesticus*, *M. musculus molossinus*, *M. musculus castaneus*, *M. spretus* (Spain), and *M. dumni* were tested with the multiplex PCR primers to determine assay robustness. Full unique profiles amplified in the designated allele range were obtained from the panel for all but the following samples: CAST/EiJ (*M. musculus castaneus*), JF1/Ms (*M. musculus molossinus*), SPRET (*M. spretus*), and *M. dumni* cell line. DNA from CAST and JF1 mice resulted in amplicons for each marker. However, the PCR product was outside of the designated allele range for the 18-3 and 6-7 loci, respectively. Sequencing the CAST mouse DNA revealed that this sample has conserved sequence flanking the repeat region. However, fifty-two ATCT repeats were observed at this locus. Thus, twenty-nine more repeats were observed at this locus than in the designated allele range.

Because of additional repeats present in the CAST mouse sample, the amplified product appears between STR markers 4-2 and 6-7. All *M. musculus molossinus* samples resulted in full profiles except for DNA from the JF1 mouse which amplified outside the designated allele range for marker 6-7. The additional thirty-two repeats that JF1 contains at the 6-7 locus may be explained in the origin of *Mus musculus molossinus*, a natural hybrid of *M. m. musculus* and *M. m. castaneus*, the latter shown to deviate from the designated allele range at marker 18-3.

DNA from the SPRET mouse (*M. spretus*) results in amplicons that fall outside the designated allele range for the following loci: 18-3, 4-2, 15-3, and X-1. The SPRET sample was sequenced at the 18-3 locus resulting in sixty-six repeats, eleven of which were GTCT repeats embedded within the defined ATCT repeat for this marker. DNA extracted from the *M. dumni* cell line does not amplify at the 6-4 STR marker and falls outside the designated allele range for X-1. Further analysis of DNA from *M. dumni* and SPRET was not continued as their profiles were incomplete using the multiplex assay. Interestingly, CAST and SPRET are mapped together in group 2 in a published mouse family tree. However, full profiles within the allele range are

observed for the other members in that group including PERC (*M. m. domesticus*), MOLG (*M. m. molossinus*), and MOLF (*M. m. molossinus*).

A panel of rodent and porcine DNA (rat, hamster, gerbil, pig), human cell lines (HeLa, HEPM, SK-BR-3, MCF10A) and nonhuman primate DNA samples (Vero, COS-7, rhesus, baboon, cynomolgus monkey) were tested with the multiplex assay to determine assay specificity. None of these samples resulted in a complete profile using the primers targeting mouse STR markers. DNA from Wistar, Fischer, and Sprague-Dawley rats resulted in a single amplified product in the red dye channel; however, each sample resulted in an amplicon with a fragment length of 219 base pairs. Characteristic stutter peaks associated with polymerase slippage of repeat regions were absent in the rat samples. Lack of stutter peaks and identical amplicon sizes for each rat strain suggests the peak present is most likely a PCR artifact rather than amplification of a repeat region. Amplification products were absent for each mouse STR marker when DNA from human and African green monkey cell lines were tested. However, both cell lines amplified at the human STR markers (D8S1106 and D4S2408) present in the multiplex as expected. No significant amplicons were visible for pig, hamster, or gerbil DNA.

SNP assays, commonly used to type mouse strains, are efficient in discriminating between different strains of mice, but may not be ideal in differentiating between cell lines derived from the same substrain. SNPs are mostly bi-allelic markers whereas STR markers typically have greater than five alleles. Using the mouse multiplex assay, unique profiles were obtained for the mouse cell lines listed in FIG. 5 with the capability of distinguishing between three Balb/c-derived cell lines. FIG. 5 is a table illustrating the complete genetic profiles of six mouse cell lines in accordance with the present disclosure.

In FIG. 4, the repeat numbers are listed for each locus. Microvariants are indicated by a decimal point. As shown in FIG. 4, there are many conserved alleles between the three Balb/c-derived samples; however, there are sufficient differences resulting in unique profiles for each individual cell line. Two of the Balb/c-derived cell lines, mouse myeloma cells (P3X63Ag8.653) and hybridoma cells (HK-PEG-1), are very similar in their genotype, only varying by one allele at the 9-2 locus. The HK-PEG-1 cell line was produced by fusing P3X63Ag8.653 (myeloma cells originating from a BALB/c mouse) with spleen cells from a BALB/c mouse, explaining why they share so many alleles. The myeloma cell line is heterozygous at the 9-2 locus whereas the hybridoma cell line is homozygous. To verify the presence of a null allele at the 9-2 marker, a panel of primers was tested with DNA from the hybridoma cells resulting in amplicons ranging from 132 to 244 base pairs (bp). Homozygote peaks were present in each sample, supporting the findings that these two cell lines differ by one allele at this marker.

To test assay sensitivity and determine the lower limits of detection, DNA from NIH3T3, HeLa, and Vero cell lines was diluted from 6 nanograms (ng) to 7.8 picograms (pg). A full profile for NIH3T3 cells was obtained using 62 pg of DNA but resulted in a loss of an allele at one mouse STR markers at 31 pg of DNA. The two human STR markers were also tested and resulted in peaks above the analytical threshold (50 relative fluorescent units) for HeLa and Vero cell lines using 62 and 187 pg of DNA, but resulted in allelic drop-out at 31 and 93 pg of DNA, respectively. In previous studies, higher concentrations of Vero cell DNA (6 ng) were needed to obtain an STR profile using human STR markers

when compared to human DNA (0.5-1 ng). This is consistent with the higher concentrations of Vero DNA needed in this study to amplify efficiently using the human STR markers in the multiplex assay.

The multiplex assay described herein was designed to detect human or African green monkey cell line contamination of mouse cells by incorporating two human STR markers that amplify outside the designated allele ranges for the nine mouse STR markers. Mixture ratios ranging from 1:1 to 10:1 of NIH3T3 and HeLa DNA were tested to model contamination scenarios.

Referring now to FIG. 5, illustrated is a genetic profile of the NIH3T3 cell line using the multiplex assay of the present disclosure. An electropherogram depicting a pure NIH3T3 STR profile is shown in this FIG. 7.

Referring now to FIG. 6, illustrated is a human contaminant detected in the NIH3T3 STR profile. A 1:1 ratio of NIH3T3 and HeLa DNA is shown in this FIG. 8. Even at the lowest dilution of HeLa DNA (90 pg), human STR markers were detected above the analytical threshold. The assay can also be used to detect a mixture of multiple mouse cell lines.

Referring now to FIG. 7, illustrated is a genetic profile of the RAW 264.7 cell line using the mouse multiplex assay (1 ng DNA). This electropherogram depicts a pure RAW264.7 STR profile. Mixture ratios ranging from 1:1 to 10:1 of NIH3T3 and RAW264.7 DNA were tested and full profiles of both cell lines were present even at the lowest DNA dilution (90 pg).

Referring now to FIG. 8, illustrated is a mixture of NIH3T3 and RAW 264.7 mouse cell lines detected using the assay described herein. FIG. 10 shows a 1:1 mixture of the two mouse cell lines.

The majority of mouse cell lines are derived from inbred mice resulting in alleles that are mostly homozygous in nature. For example, as shown earlier in connection with FIG. 4, the RAW 264.7 mouse cell line is homozygous at each STR marker. Multiple alleles present at each locus could indicate a mixed population of cells. Triallelic patterns have been observed in some human cell lines at a particular locus, which may or may not be equal in intensity. The L929 cell line appears to have three alleles with similar peak height intensities at the 15-3 marker and each allele is four base pairs or one repeat apart. Since most of the mouse samples tested were homozygous for the majority of the markers, a panel of primers targeting the 15-3 locus were tested in monoplex with DNA from L929 cells. The amplicons ranged from 210 to 435 base pairs in length and each resulted in three alleles that were four bases apart with very little peak height imbalance. The evidence supports a true triallelic pattern at the 15-3 marker.

Alteration of genetic profiles of some cancer cell lines has been observed previously at high passage numbers. However, other studies show STR stability over high passage numbers in some human cancer cell lines and in African green monkey cell lines. Accordingly, stability may be cell line dependent. To test the stability of the mouse STR markers in this assay, L929 and NIH3T3 cell lines were carried independently and in duplicate flasks up to passage 44 and 45, respectively.

Genotypes were determined and standard deviations were calculated for each locus representing the variations in fragment lengths over all passage numbers. The NIH3T3 cell line resulted in the lowest standard deviation values (0.02-0.05) for each locus. The L929 cell line resulted in standard deviations ranging from 0.05 to 0.14. The STR markers with the highest standard deviations in L929 cells are 6-7 (0.14) and 5-5 (0.13). In both the NIH3T3 and L929

cell lines, even the highest standard deviation values did not result in an allele repeat number change indicating stable STR profiles at high passage numbers. The changes in fragment lengths for each marker over the passage period were not significant enough to change the allele calls and the variability in the amplicon sizes fell within the range of the instrument fluctuation. Identical DNA samples were tested on three different days using the same instrument and the variation in fragment length was  $\pm 0.3$  base pairs.

In addition to stability of the STR profile for NIH3T3 cells over time, profile stability was evaluated after transfection procedures. The TN1 cell line, derived from NIH3T3 cells obtained from ATCC in 2003 and engineered to express the gene for green fluorescent protein, was analyzed using the multiplex assay and resulted in identical STR profiles for both TN1 and recently obtained NIH3T3 cells. These data support the findings that the STR markers are stable over time in transfected NIH3T3 cell lines.

The mouse multiplex assay described herein can be used to identify cell lines derived from *M. musculus musculus* and *M. musculus domesticus* species. The assay is also useful in identifying *M. musculus molossinus* and *M. musculus castaneus* species which amplify at each locus, but in some instances failed to fall within the designated allele range for one of the STR markers. This assay may not be suitable for genotyping mouse cell lines derived from *M. spretus* (amplicons may fall outside the designated allele range for four STR markers) or *M. dummi* which may fail to amplify at the 6-4 locus.

Stability studies show the mouse STR markers are stable with high passage numbers and the STR profiles remain unchanged after transfection procedures in the TN1 cell line. Although the STR markers are stable up through passages 44-45, it may be desirable to genotype samples at low passage numbers. The power of discrimination based on the probability of a random match is 1 in 5.7 million using the nine STR markers in the multiplex assay. The assay described in the present disclosure can be used to identify both human and African green monkey cell line contaminants using the two human STR markers incorporated in the multiplex assay in addition to detecting mixtures of mouse cell lines. The targeted tetranucleotide repeat regions in the mouse genome result in unique individual profiles making this assay more sensitive and specific than those that are currently available. The requirement of cell line authentication is becoming more routine, and this assay provides a reliable method to genotype mouse cell lines.

The STR profiles shown in the Table in FIG. 9 display the allele range for the 9 markers. This data could be used to develop a size reference standard that is a locus-specific allelic ladder.

FIG. 10 is another table illustrating primers for STR amplification and their corresponding chromosomal locations in accordance with one embodiment of the present disclosure. In FIG. 10, mouse chromosomal locations, or base pairs (bp), are based on the GRCh37.p13 Primary Assembly mouse build. Primer concentrations in the right-most column are final concentrations of forward and reverse primers in a 20 reaction volume except for primers 2-1 and 13-1 which were 0.100  $\mu\text{M}$  and 0.250  $\mu\text{M}$  respectively. Primer concentrations were determined empirically based on peak height, DNA concentration and the number of cycles in the PCR program.

Based on FIG. 10, the exemplary amplification primers are set forth below. In order to specifically amplify the selected STR markers, the following oligonucleotide primer pairs were used, where "F" and "R" correspond to the

forward and reverse primers respectively. Additional numbers found with these "F" and "R" designations, e.g., R4, F4, R7, are simply used to keep track of the different forward and reverse primers. The PCR is the product expected from amplification of the particular STR locus to which the oligonucleotide pair was directed.

Mo11-1  
 F4: (SEQ ID NO: 34) [6FAM]-CCAGCCAGCCACTCTAGTTA  
 R4: (SEQ ID NO: 35) GTTTCTTACAGCTTCAGGACCATGGAA  
 Primer (Seq. ID No. 97)  
 Mo19-2  
 F4: (SEQ ID NO: 36) [6FAM]-AGGCTAGCACTGTTCCCTTGT  
 R7: (SEQ ID NO: 37) GTTTCTTACTCAGCACCTTCCATCCTG  
 Primer (Seq. ID No. 98)  
 Mo1-2  
 F3: (SEQ ID NO: 38) [VIC]-TCTTTAAAAATCAAACAGGCAAA  
 R: (SEQ ID NO: 39) GTTTCTTGGGGAGGTGGGGGTGTATAA  
 Primer (Seq. ID No. 99)  
 Mo7-1  
 F: (SEQ ID NO: 40) [VIC]-ACACCAAGTTGGGGACTAGGA  
 R: (SEQ ID NO: 41) GTTTCTTAACTGGTCTGGGTCTGAAC  
 Primer (Seq. ID No. 100)  
 Mo1-1  
 F: (SEQ ID NO: 42) [VIC]-CCCTTCACTCCTTCATTCCA  
 R: (SEQ ID NO: 43) GTTTCTTTGAGCCTAAGGACCTGGACA  
 Primer (Seq. ID No. 101)  
 Mo3-2  
 F3: (SEQ ID NO: 44) [VIC]-TGAGCTACCATGTGGGTACA  
 R3: (SEQ ID NO: 45) GTTTCTTCACACACACACAAAGATGGA

-continued

Primer (Seq. ID No. 102)  
 5 Mo8-1 (SEQ ID NO: 46)  
 F4: [VIC]-AGTAATATCCTGGTCCTGGCC  
 R4: (SEQ ID NO: 47) GTTTCTTGAGCTCACTATGTAGCTATTGGA  
 Primer (Seq. ID No. 103)  
 10 Mo2-1 (SEQ ID NO: 48)  
 F2: [NED]-ACAACTTCAAACCTTGCTGGAA  
 R3: (SEQ ID NO: 49) GTTTCTTTCCTGCTTAGCTTGTCAATTTCC  
 Primer (Seq. ID No. 104)  
 20 Mo11-2 (SEQ ID NO: 50)  
 F2: [NED]-AAGGCAGGGGAATTCACAGT  
 R4: (SEQ ID NO: 51) GTTTCTTCTCACCATTGCAGTCCTGA  
 Primer (Seq. ID No. 105)  
 30 Mo17-2 (SEQ ID NO: 52)  
 F: [PET]-CTCTTCTCCATCCGTCCATC  
 R: (SEQ ID NO: 53) GTTTCTTATGGCTCTTGCCACAAATCT  
 Primer (Seq. ID No. 106)  
 40 Mo13-1 (SEQ ID NO: 54)  
 F4: [PET]-AACTCCCACTGCTTCTTGGT  
 R6: (SEQ ID NO: 55) GTTTCTTGCCAAAGTAAACCCCTTCTCCC  
 Primer (Seq. ID No. 107)  
 50  
 55  
 60  
 65

A pigtail sequence "X" may be added to the 5' end of the reverse primer. The pigtail sequence must contain a guanine (G). The pigtail sequence may contain a G and a cytosine (C). The pigtail sequence may contain a G and thymidine (T). The pigtail sequence may contain a G, C, and T. For example, a suitable pigtail sequence/reverse primer for Mo 13-1 is as follows: R primer (R6pig):

GTTCCTTGCCAAAGTAAACCCCTTCTCCC

In order to implement the method and/or use a kit for determining the alleles present in a DNA sample, a DNA sample may be obtained for analysis. Then, selection may occur for a set of STR marker loci of the DNA sample to be analyzed that can be amplified together in a multiplex

amplification reaction. The source of the DNA sample to be analyzed is at least one of a mouse and a cell line derived from a mouse. The set of STR marker loci are selected from the group consisting of: Mo11-1, Mo19-2, Mo1-2, Mo7-1, Mo1-1, Mo3-2, Mo8-1, Mo2-1, Mo11-2, Mo17-2, Mo13-1.

Then, a set of oligonucleotide primer pairs may be selected. Each oligonucleotide primer pair in the set flanks a single locus in the set of STR marker loci. Each oligonucleotide primer pair is capable of amplification of a single locus from the set of STR marker loci in a multiplex amplification reaction. Then, the set of STR marker loci is co-amplified in a multiplex amplification reaction. The product of the multiplex amplification reaction comprises a mixture of amplified alleles from each of the co-amplified loci in the set of STR marker loci. Then, evaluation of the products of the co-amplification reaction may occur in order to determine the alleles present at each of the loci analyzed in the set of STR marker loci within the DNA sample.

FIG. 11 is another table illustrating sequencing primers and their respective amplicon size range as well as annealing temperatures in accordance with one embodiment of the present disclosure.

As illustrated in FIG. 11, the STR marker, forward and reverse primers, their respective amplicon size range as well as annealing temperatures, respectively, are as follows:

Mo11-1	FSeq1: CACCCATGAAAGAGCCAAGT* (SEQ ID NO: 60) RSeq1: TTCAGGACCATGGAAAGCTC* (SEQ ID NO: 61)	342-367	60
Mo19-2	FSeq1: TCCTCAAGGCTAGCACTGTTC* (SEQ ID NO: 62) RSeq1: AAATCCAGCACTCAGGAGA* (SEQ ID NO: 63)	276-300	60
Mo1-2	F: ACTCCTCCCAGATTCCCTTCG* (SEQ ID NO: 64) RSeq2: TGCCCATATGTATGGTTCC* (SEQ ID NO: 65)	199-236	60
Mo7-1	F: ACACCAGTTGGGGACTAGGA (SEQ ID NO: 66) R3: CTGCTCTTCCAGAGGTCCTG* (SEQ ID NO: 67)	349-405	60
Mo1-1	F: CCCTTCACTCCTTCATTCCA (SEQ ID NO: 68) R: TGAGCCTAAGGACCTGGACA (SEQ ID NO: 69)	325-366	59
Mo3-2	F: CCATGTGGGTACATTGTTAAGC* (SEQ ID NO: 70) R: CATGTGATAGAGGATGGATAGATGA* (SEQ ID NO: 71)	277-347	59
Mo8-1	F: TGGCAGAGTGCATACCCTAGC* (SEQ ID NO: 72) R: CTAGAACTGGGCCTGGTGGT* (SEQ ID NO: 73)	271-325	59
Mo2-1	FSeq1: CCTAGAGTCTTTGTGGCTGGA* (SEQ ID NO: 74) RSeq1: CACCTAGGTCTCCGATTGA* (SEQ ID NO: 75)	373-408	60
Mo11-2	F: TCCAGGGGAATCTTTGACAC* (SEQ ID NO: 76) R: GAGGGCTGGTAAAGTTGTGC* (SEQ ID NO: 77)	203-260	59

-continued

Mo17-2	F: CTCTTCTCCATCCGTCCATC (SEQ ID NO: 78) R: ATGGCTCTGGCCAAAATCT (SEQ ID NO: 79)	183-227	59
Mo13-1	F: TCGTTACCATGCCTCTTTCC* (SEQ ID NO: 80) R3: CGCCACAGCCTCATTACTC* (SEQ ID NO: 81)	309-345	60

FIG. 12 is a table defining STR fragment length and corresponding repeat number in accordance with one embodiment of the present disclosure.

A kit may be provided for determining the alleles present in a DNA sample to be analyzed. The kit may include, among other things, oligonucleotide primer pairs for co-amplifying a set of STR marker loci of at least one DNA sample to be analyzed. The set of loci may comprise one or more of STR loci selected from the new group consisting of STR marker loci Mo11-1, Mo19-2, Mo1-2, Mo7-1, Mo1-1, Mo3-2, Mo8-1, Mo2-1, Mo11-2, Mo17-2, Mo13-1, and/or the old and new groups of STR marker loci combined: 18-3, 4-2, 6-7, 15-3, 6-4, 12-1, 5-5, X-1, 11-1, 19-2, 1-2, Mo7-1, Mo1-1, Mo3-2, Mo8-1, Mo2-1, Mo11-2, Mo17-2, Mo13-1, D8S1106 and D4S2408.

A size standard may be used to run the multiplex assay which is commercially available (e.g., ABI GeneScan 600 LIZ size standard). The kit may have an allelic ladder, or a size standard included. Where a size reference standard is used, the size reference standard may comprise a locus-specific allelic DNA size marker.

The size reference standard may comprise a DNA marker of known size. The DNA marker could be labeled with a label distinguishable from the label attached to the oligonucleotide primers.

The allelic ladder (different than a size standard) may have the same dye as the corresponding allele. For example, for Mo11-1 which is labeled with FAM, the alleles for that marker may also be amplified with a FAM dye so that the fragments may migrate the same. For the allelic ladder, the label or dye will correlate with the corresponding STR marker for those alleles. For example, 18-3 alleles may be labeled with FAM so they migrate together. By contrast, the sizing standard—which is commercially available—such as the GeneScan LIZ600, has a different dye distinguishable from the STR markers.

Four (4) of the primer sequences illustrated and discussed hereinabove, have been updated. The four (4) updated primer sequences listed in FIG. 13 are as follows: (1) D8S1106 (human); (2) 15-3 (mouse); (3) 12-1 (mouse) and X-1 (mouse). The new 15-3, 12-1, X-1 are from GRCh38.p4 mouse build and the D8S1106 is from the human GRCh37.p13 build.

These four primers may be used in the multiplex assay. They are not going to replace the primers in FIG. 1, but may be used in combination with the eleven (11) new STR markers of FIG. 10 in an updated multiplex assay. These new four primer sequences needed to be changed in order to fit these eleven (11) new STR markers into the new assay. It may be desirable for each STR marker to have twenty (20) to thirty (30) base pairs in between each other so that there is no overlap.

Referring now to FIG. 13, illustrated are updated primers for the nine (9) mouse STR markers of FIG. 1, in accordance with one embodiment of the present disclosure. For some of these updated primers in FIG. 13, they will also work with the new eleven (11) markers listed in FIG. 10 hereinabove.

## 21

These updated primers sequences allow for the additional STR markers of FIG. 10 to be added to the multiplex assay. For each of these updated primers the STR marker region will be the same. However, the amplicon or product may vary (longer or shorter) in order to provide for optimal spacing between each STR marker.

Regarding updated primer sequence D8S1106, the primer sequence is different from that listed earlier in connection with FIG. 1. However the STR repeat region is the same. This updated primer sequence for D8S1106 can be used to screen for contamination of mouse cell lines with human or African green monkey cell lines. This updated primer sequence for D8S1106 can also be used to identify human and African green monkey cell lines (e.g., Vero cells). The chromosomal locations for human STR markers D8S1106 are based on the GRCh37.p13 Primary Assembly (*Homo sapiens* chromosome 8 genomic contig.), an updated version of the NCBI 37.3 human build.

The forward/reverse primer pair for new updated primer sequence D8S1106 is as follows:

F2 Primer: (Seq. ID No. 108)  
TCTGCTGATTGTGCCCATG

R3 Primer: (Seq. ID No. 109)  
ATTGGCTGGTTAGTGACACA

Regarding the new updated primer sequence for "15-3", this primer sequence refers to the STR marker on mouse chromosome 15, corresponding to base pairs 4930200-4930500 (NCBI 38.1 mouse build, GenBank Accession # NT\_039617.8, >gi|149266402:1880200-1880500 *Mus musculus* strain C57BL/6J chromosome 15 genomic contig, MGSCv37 C57BL/6J). The forward/reverse primer pair for new updated primer sequence 15-3 is as follows:

F- (SEQ ID NO: 87)  
TCTGGCGTGTCTGTCATAA

R-E- (SEQ ID NO: 88)  
GTTTCTTGAGGCTATCTAACTTGACATGCT

Please note that the GTTTCTT sequence is underlined here and elsewhere to indicate that this sequence is not found in the genome but added for adenylation issues. Based on the GRCm38.4 build, GenBank Accession # NC\_000081.6 (corresponding base pairs now 4930233-4930450). The underlined sequence above is not part of genome sequence. It is added to stabilize the primers and promotes full adenylation of products during PCR.

As earlier described in connection with primer sequence 15-3, the L929 cell line can have three alleles with similar peak height intensities at the 15-3 marker and each allele may be four base pairs or one repeat apart. The amplicons may result in three alleles that are four bases apart with very little peak height imbalance. The evidence supports a true triallelic pattern at the 15-3 marker.

## 22

Regarding updated primer sequence 12-1, the primers have changed. The forward and reverse primers are as follows:

F2: (SEQ ID NO: 89)  
TTTCAAATGTCATTGAACACA

R6: (SEQ ID NO: 90)  
GTTTCTTTGGTCCTTCAGTATCATCCTTG

The underlined sequence is not part of the genome sequence. It is added to stabilize the primers and allow for full adenylation of products during PCR. This "12-1" still refers to the STR marker on mouse chromosome 12. However, the updated base pairs are now located at GRCm38.p4 NC\_000078.6 (38480946-38481212). It's an updated mouse build 38.p4. The 12-1 primers listed below will only work with the original 9 STR markers of FIG. 1. The primers for 12-1 were redesigned to allow better spacing between STR markers 12-1 and 17-2. If the markers are too close together, the genotypes will be impossible to determine as it will be difficult to distinguish which allele goes with which marker. These 12-1 primers will overlap the allele range for the new 17-2 STR marker, which may cause issues in determining which allele goes with which STR marker.

Updated primer X-1 differs from the earlier one we referenced hereinabove in that the forward primer is updated, but the reverse primer remains the same. The forward/reverse primers for updated X-1 primer sequence are as follows:

F: (SEQ ID NO: 93)  
AACAAAAATGTCCTCAATGC

R: (SEQ ID NO: 94)  
GTTTCTTAAGGTATATATCAAGATGGCATTATCA

Regarding the new updated primer sequence for X-1, this "X-1" refers to the STR marker on mouse chromosome X, corresponding to base pairs 55739301-55739725 (NCBI 38.1 mouse build, GenBank Accession # NT\_039706.7, >gi|149271867:55739107-55739854 *Mus musculus* strain C57BL/6J chromosome X genomic contig, MGSCv37 C57BL/6J).

Referring now to FIG. 14, illustrated are the four new primers and allele distribution of changes to eight of the original nine (9) STR markers (to accommodate these primers into the new twenty-one-plex (21-plex) assay). The 21-plex assay may include STR marker loci 18-3, 4-2, 6-7, 15-3, 6-4, 12-1, 5-5, X-1, Mo11-1, Mo19-2, Mo1-2, Mo1-1, Mo1-1, Mo3-2, Mo8-1, Mot-1, Mo11-2, Mo17-2, Mo13-1, D8S1106 and D4S2408.

While the specification describes particular embodiments of the present invention, those of ordinary skill can devise variations of the present invention without departing from the inventive concept.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 102

<210> SEQ ID NO 1





-continued

---

```

<210> SEQ ID NO 6
<211> LENGTH: 294
<212> TYPE: DNA
<213> ORGANISM: Mouse

<400> SEQUENCE: 6
tttgcaacag ctcagtttcc ttgagaaaat tgctttaata caaatcccta ggaatctaca    60
tttctoctag gaaaggtaat tttttagata gatagatgga gatagataga tagatagata    120
gatagataga tagatagata gatagataga tagatagatg atagatagat aaattctact    180
ggatagaaag atatcatgta ggctgagata ctacctactt atagtgaagt agtgaaggaa    240
attcataagc ggtaaatata aagtattgga actacctaag atctgccagc gatt          294

```

```

<210> SEQ ID NO 7
<211> LENGTH: 224
<212> TYPE: DNA
<213> ORGANISM: Mouse

<400> SEQUENCE: 7
caaaattgtc attgaacaca tgtaaaaaag cagaagatag atagatagat agatagatag    60
atagatagat agatagatag atagatagat gatagataga tagatattatc taggagcata    120
gatgtaataa ctaaattgta taatgtgaca ttaaataagt ccatcactat aaaataatct    180
ttaataatta caaaaagttg tacttcagta tttcttgacc attg                    224

```

```

<210> SEQ ID NO 8
<211> LENGTH: 341
<212> TYPE: DNA
<213> ORGANISM: Mouse

<400> SEQUENCE: 8
cgttttacct ggctgacaca cttcctgggc cgggtggctcc tggttttgaa agaaatcctt    60
gatggatgtc ataataaaga ggcttggtgg tgcggagcgc gaggatccgg gttccggttt    120
ctgcctttat ctatctatct atctatctat ctatctatct atctatctat ctatctatct    180
atctatctat ctatctatta tttatttatt caggaaatgc ttacttaggg agctgggtag    240
caacaagggt tttctgtac ctgctaggaa caggcaagca tccctaagtc agtttgtagt    300
ctttattaag caataaattg tttggtattg agttttaaac c                        341

```

```

<210> SEQ ID NO 9
<211> LENGTH: 402
<212> TYPE: DNA
<213> ORGANISM: Mouse

<400> SEQUENCE: 9
ggatggatgg atggatgaaa agaatagata gatagataga tagatagata gatagataga    60
tagatagata gatagataga tgatagatag atagatagat agatagatag atagatagat    120
agatagatag aaagaatgta atacatatga gcatactact cagttttaaa aattaatttc    180
tgccatttga tattattgac ctagaggaaa atatgttaca ttaaataacc tgtggaaatg    240
agtaggatac cacacaatcc catttaaatg ttttatatac atcttcactg ctcccgtgt    300
gacaaatagt gtccttattt ttgaatgttc taaaatattt tggtgggttaa tattcattgt    360
acatagtcac gaatatgata atgccatctt gatataatcc tt                        402

```

```

<210> SEQ ID NO 10

```

-continued

---

<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Mouse

<400> SEQUENCE: 10

tctttctcct ttgtgtcatg c 21

<210> SEQ ID NO 11  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Mouse

<400> SEQUENCE: 11

gtttcttgct aaataactaa gcaagtgaac aga 33

<210> SEQ ID NO 12  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Mouse

<400> SEQUENCE: 12

aagcttctct ggccatttga 20

<210> SEQ ID NO 13  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Mouse

<400> SEQUENCE: 13

gttcataaac ttcaagcaat gaca 24

<210> SEQ ID NO 14  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Mouse

<400> SEQUENCE: 14

agtcacccca gtgcattctc 20

<210> SEQ ID NO 15  
<211> LENGTH: 27  
<212> TYPE: DNA  
<213> ORGANISM: Mouse

<400> SEQUENCE: 15

gtttcttcat gtggctggta tgctgtt 27

<210> SEQ ID NO 16  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Mouse

<400> SEQUENCE: 16

ggattgcaa gaatttgagg 20

<210> SEQ ID NO 17  
<211> LENGTH: 29  
<212> TYPE: DNA  
<213> ORGANISM: Mouse

<400> SEQUENCE: 17

gtttctttcc tgagttgtgg acagggtta 29

-continued

---

<210> SEQ ID NO 18  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Mouse  
  
 <400> SEQUENCE: 18  
  
 tctgggcgtg tctgtcataa 20

<210> SEQ ID NO 19  
 <211> LENGTH: 27  
 <212> TYPE: DNA  
 <213> ORGANISM: Mouse  
  
 <400> SEQUENCE: 19  
  
 gtttcttttc tcagggagga gtgtgct 27

<210> SEQ ID NO 20  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Mouse  
  
 <400> SEQUENCE: 20  
  
 tttgcaacag ctcagtttcc 20

<210> SEQ ID NO 21  
 <211> LENGTH: 27  
 <212> TYPE: DNA  
 <213> ORGANISM: Mouse  
  
 <400> SEQUENCE: 21  
  
 gtttcttaat cgctggcaga tcttagg 27

<210> SEQ ID NO 22  
 <211> LENGTH: 25  
 <212> TYPE: DNA  
 <213> ORGANISM: Mouse  
  
 <400> SEQUENCE: 22  
  
 caaaattgtc attgaacaca tgtaa 25

<210> SEQ ID NO 23  
 <211> LENGTH: 35  
 <212> TYPE: DNA  
 <213> ORGANISM: Mouse  
  
 <400> SEQUENCE: 23  
  
 gtttctttca atggtcaaga aatactgaag tacia 35

<210> SEQ ID NO 24  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Mouse  
  
 <400> SEQUENCE: 24  
  
 cgttttacct ggctgacaca 20

<210> SEQ ID NO 25  
 <211> LENGTH: 31  
 <212> TYPE: DNA  
 <213> ORGANISM: Mouse  
  
 <400> SEQUENCE: 25  
  
 gtttcttggg ttaaaactca ataccaaca a 31

-continued

---

<210> SEQ ID NO 26  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Mouse  
  
 <400> SEQUENCE: 26  
  
 ggatggatgg atggatgaaa 20

<210> SEQ ID NO 27  
 <211> LENGTH: 34  
 <212> TYPE: DNA  
 <213> ORGANISM: Mouse  
  
 <400> SEQUENCE: 27  
  
 gtttcttaag gtatatatca agatggcatt atca 34

<210> SEQ ID NO 28  
 <211> LENGTH: 13  
 <212> TYPE: DNA  
 <213> ORGANISM: mouse  
  
 <400> SEQUENCE: 28  
  
 attggtcaat atg 13

<210> SEQ ID NO 29  
 <211> LENGTH: 12  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus  
  
 <400> SEQUENCE: 29  
  
 gtaccactga ga 12

<210> SEQ ID NO 30  
 <211> LENGTH: 19  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
  
 <400> SEQUENCE: 30  
  
 gtttaccct gcatcatgg 19

<210> SEQ ID NO 31  
 <211> LENGTH: 30  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
  
 <400> SEQUENCE: 31  
  
 gtttctttca gaattgctca tagtgcaaga 30

<210> SEQ ID NO 32  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
  
 <400> SEQUENCE: 32  
  
 tcatttccat agggtaagtg aaaa 24

<210> SEQ ID NO 33  
 <211> LENGTH: 27  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
  
 <400> SEQUENCE: 33  
  
 gtttcttgcc atggggataa aatcaga 27

-continued

---

<210> SEQ ID NO 34  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus  
  
 <400> SEQUENCE: 34  
 ccagccagcc actctagtta 20

<210> SEQ ID NO 35  
 <211> LENGTH: 27  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus  
  
 <400> SEQUENCE: 35  
 gtttcttaca gcttcaggac catggaa 27

<210> SEQ ID NO 36  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus  
  
 <400> SEQUENCE: 36  
 aggctagcac tggtccttgt 20

<210> SEQ ID NO 37  
 <211> LENGTH: 27  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus  
  
 <400> SEQUENCE: 37  
 gtttcttact cagcaccttc catcctg 27

<210> SEQ ID NO 38  
 <211> LENGTH: 23  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus  
  
 <400> SEQUENCE: 38  
 tctttaaaaa tcaaacaggc aaa 23

<210> SEQ ID NO 39  
 <211> LENGTH: 27  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus  
  
 <400> SEQUENCE: 39  
 gtttcttggg gaggttgggg tgtataa 27

<210> SEQ ID NO 40  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus  
  
 <400> SEQUENCE: 40  
 acaccagttg gggactagga 20

<210> SEQ ID NO 41  
 <211> LENGTH: 27  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus  
  
 <400> SEQUENCE: 41

-continued

---

gtttcttaac tgggtctggg tctgaac	27
<210> SEQ ID NO 42 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Mus musculus <400> SEQUENCE: 42	
cccttcactc cttcattcca	20
<210> SEQ ID NO 43 <211> LENGTH: 27 <212> TYPE: DNA <213> ORGANISM: Mus musculus <400> SEQUENCE: 43	
gtttctttga gcctaaggac ctggaca	27
<210> SEQ ID NO 44 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Mus musculus <400> SEQUENCE: 44	
tgagctacca tgtgggtaca	20
<210> SEQ ID NO 45 <211> LENGTH: 29 <212> TYPE: DNA <213> ORGANISM: Mus musculus <400> SEQUENCE: 45	
gtttcttcac acacacacac aaagatgga	29
<210> SEQ ID NO 46 <211> LENGTH: 21 <212> TYPE: DNA <213> ORGANISM: Mus musculus <400> SEQUENCE: 46	
agtaatatcc tggctctggc c	21
<210> SEQ ID NO 47 <211> LENGTH: 30 <212> TYPE: DNA <213> ORGANISM: Mus musculus <400> SEQUENCE: 47	
gtttcttgag ctcactatgt agctattgga	30
<210> SEQ ID NO 48 <211> LENGTH: 22 <212> TYPE: DNA <213> ORGANISM: Mus musculus <400> SEQUENCE: 48	
acaacttcaa actttgctgg aa	22
<210> SEQ ID NO 49 <211> LENGTH: 29 <212> TYPE: DNA <213> ORGANISM: Mus musculus <400> SEQUENCE: 49	

-continued

---

gtttctttcc tgettagctt gtcatttcc	29
<210> SEQ ID NO 50 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Mus musculus <400> SEQUENCE: 50	
aaggcagggg aattcacagt	20
<210> SEQ ID NO 51 <211> LENGTH: 27 <212> TYPE: DNA <213> ORGANISM: Mus musculus <400> SEQUENCE: 51	
gtttctttct caccattgca gtctctga	27
<210> SEQ ID NO 52 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Mus musculus <400> SEQUENCE: 52	
ctcttctcca tccgtccatc	20
<210> SEQ ID NO 53 <211> LENGTH: 27 <212> TYPE: DNA <213> ORGANISM: Mus musculus <400> SEQUENCE: 53	
gtttcttatg gctcttgcca caaatct	27
<210> SEQ ID NO 54 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Mus musculus <400> SEQUENCE: 54	
aactcccact gcttcttgg	20
<210> SEQ ID NO 55 <211> LENGTH: 28 <212> TYPE: DNA <213> ORGANISM: Mus musculus <400> SEQUENCE: 55	
gtttcttgcc aaagtaaacc cttctccc	28
<210> SEQ ID NO 56 <211> LENGTH: 21 <212> TYPE: DNA <213> ORGANISM: Mus musculus <400> SEQUENCE: 56	
aaatgttagg ggtagggagc a	21
<210> SEQ ID NO 57 <211> LENGTH: 27 <212> TYPE: DNA <213> ORGANISM: Mus musculus	

-continued

---

<400> SEQUENCE: 57  
gtttctttaa gttccaggcc atccaag 27

<210> SEQ ID NO 58  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 58  
ttggtgtctt atggcttcca ga 22

<210> SEQ ID NO 59  
<211> LENGTH: 27  
<212> TYPE: DNA  
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 59  
gtttctttgc cattttccac acacact 27

<210> SEQ ID NO 60  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 60  
cacccatgaa agagccaagt 20

<210> SEQ ID NO 61  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 61  
ttcaggacca tggaaagctc 20

<210> SEQ ID NO 62  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 62  
tctctcaaggc tagcactgtt c 21

<210> SEQ ID NO 63  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 63  
aaatcccagc actcaggaga 20

<210> SEQ ID NO 64  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 64  
actctctcca gattccttcg 20

<210> SEQ ID NO 65  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Mus musculus



-continued

---

<400> SEQUENCE: 65  
tgcccatatt gtatggttc c 21

<210> SEQ ID NO 66  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 66  
acaccagtgtg gggactagga 20

<210> SEQ ID NO 67  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 67  
ctgctcttcc agaggtcctg 20

<210> SEQ ID NO 68  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 68  
cccttcactc cttcattcca 20

<210> SEQ ID NO 69  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 69  
tgagcctaag gacctggaca 20

<210> SEQ ID NO 70  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 70  
ccatgtgggt acattgttaa gc 22

<210> SEQ ID NO 71  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 71  
catgtgatag aggatggata gatga 25

<210> SEQ ID NO 72  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 72  
tggcagagtg catacctagc 20

<210> SEQ ID NO 73  
<211> LENGTH: 20  
<212> TYPE: DNA

-continued

---

<213> ORGANISM: Mus musculus  
 <400> SEQUENCE: 73  
 ctagaactgg gcctggtggt 20

<210> SEQ ID NO 74  
 <211> LENGTH: 21  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus  
 <400> SEQUENCE: 74  
 cctagagtct ttgtggctgg a 21

<210> SEQ ID NO 75  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus  
 <400> SEQUENCE: 75  
 cacctaggtc ctccgattga 20

<210> SEQ ID NO 76  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus  
 <400> SEQUENCE: 76  
 tccaggggaa tctttgacac 20

<210> SEQ ID NO 77  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus  
 <400> SEQUENCE: 77  
 gagggctggt aaagttgtgc 20

<210> SEQ ID NO 78  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus  
 <400> SEQUENCE: 78  
 ctcttctcca tccgtccatc 20

<210> SEQ ID NO 79  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus  
 <400> SEQUENCE: 79  
 atggctcttg ccacaaatct 20

<210> SEQ ID NO 80  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus  
 <400> SEQUENCE: 80  
 tcggtaccat gcctctttcc 20

<210> SEQ ID NO 81  
 <211> LENGTH: 19

-continued

---

<212> TYPE: DNA  
 <213> ORGANISM: Mus musculus  
  
 <400> SEQUENCE: 81  
  
 cgccacagcc tcattactc 19  
  
 <210> SEQ ID NO 82  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus  
  
 <400> SEQUENCE: 82  
  
 tctgggcgtg tctgtcataa 20  
  
 <210> SEQ ID NO 83  
 <211> LENGTH: 23  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus  
  
 <400> SEQUENCE: 83  
  
 aggetatcta aacttgacat gct 23  
  
 <210> SEQ ID NO 84  
 <211> LENGTH: 23  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus  
  
 <400> SEQUENCE: 84  
  
 tttcaaaatt gtcattgaac aca 23  
  
 <210> SEQ ID NO 85  
 <211> LENGTH: 22  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus  
  
 <400> SEQUENCE: 85  
  
 tggtccttca gtatcatcct tg 22  
  
 <210> SEQ ID NO 86  
 <211> LENGTH: 25  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus  
  
 <400> SEQUENCE: 86  
  
 caaaattgtc attgaacaca tgtaa 25  
  
 <210> SEQ ID NO 87  
 <211> LENGTH: 28  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus  
  
 <400> SEQUENCE: 87  
  
 tcaatggtca agaaatactg aagtacaa 28  
  
 <210> SEQ ID NO 88  
 <211> LENGTH: 21  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus  
  
 <400> SEQUENCE: 88  
  
 aacaaaaatg tccctcaatg c 21  
  
 <210> SEQ ID NO 89

-continued

<211> LENGTH: 27  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 89

aaggtatata tcaagatggc attatca 27

<210> SEQ ID NO 90  
 <211> LENGTH: 480  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 90

aactegaatt cagatccctt gcacccatga aagagccaag tgagttgggtg tgcacctgta 60  
 tcccacaata gagagacaga cacagatgga tcccctgggc ttgccagcca gccactctag 120  
 ttaaagcagt gagcttgggg tgcagtaaga gccattgcct ccaaaagtaa agtgtagaga 180  
 aattaatatg ccaactatac atacatacat acatacatac atacatacat acatacatac 240  
 atacatacat acatacatac ggggggggag agagagagaa tgagagaatg agcgagagaa 300  
 tctgaccagg acttccagac aatagagcct ttcataattt ggaaaggtgc ccttggagct 360  
 ttccatggtc ctgaagctgt taaccacagc tccattcatt tctgttcctt ccttgaataa 420  
 ggaagtgttg acctgcttgt gtgatagga agctccctgg aaacagtgt tacttaacaa 480

<210> SEQ ID NO 91  
 <211> LENGTH: 935  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 91

gctaccatgc agtgcgggg acctgaactc aagacctcca aaacagcagc caatgctccc 60  
 aaccactgag ccatccctcc agccccccct ttttttttaa ataaaggat ttaactaat 120  
 gaactatccc ctctctctt ggtgatgcta taacagaaaa ctcttagcaa agagttaagc 180  
 cttacatgta ccagtctca aggctagcac tgttctctgt ttgtagagaa attgaaacaa 240  
 agaattgacg tggcatgccc aaggtcactc agctagtaaa caggatttaa acttgtttgg 300  
 tctggttagt tccctgggaa cctcatgtct gtctgtctgt cttctatcta tcatctatgc 360  
 atctgtctat ctatctatct atctatctat ctatctatct atctatctat ctatcatcga 420  
 cactgtgctt ctctgttctc tgtgtagccc tggttgacct ggaacttgcct ctgtagagca 480  
 ggccagcttc aaaaccatag aatgccacct gtctctgtct cctgagtgct gggatttaag 540  
 gagtgtgcca gtactgccag gctgggaacc tcacttttta aatccatttt ccccctttga 600  
 tacctttttg ttgaaggaa cagactatgt cacttcacaa tgtgcttctg taacataaac 660  
 cgtaccactg gagcagggca tgacagtgca gagctggact cagtgctgtg agcaccagga 720  
 tggaaaggtgc tgagtgaagc tagcctgggc tgcatagggg aactgaaaga ttttcgaatt 780  
 cccggcccat gtacaaaatg aaagacctc gagggaagac cctcactcag aactcaagt 840  
 cccgggacag ccgctgacct aagaagacac tgagaccata cataaatgt agaaggcaag 900  
 atttaataaa gcagcgacat gaaagcacac agacc 935

<210> SEQ ID NO 92  
 <211> LENGTH: 518  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 92

-continued

---

```

ctaactccag caccactoct tcaattcct gcagctggct cctttttgce tgttttctaa    60
ttcgtgtcaa ttgtattcca ccttttttat tgaaatattt ttttcataca atatattctg    120
atcatgctct cttttttccta actcctoccca gattcctctcg atatctccaa ctcccgaatt    180
ctacagtctt tctgtcttta aaaatcaaac aggcaaattt aataaataga tagatagatg    240
atagatagat agatagatag atagatagat agatagatag atagatagat agataaaatt    300
atacaccoca acctccocccc caaagtoccca atattggaaa ccatacaata tggggcaaaag    360
gtaaaaatat caaaaacaag ctatgtgaaa caaaaagtct acaaaaatct caaaatata    420
atacaatgag tttgttttaa tttggccatc tagagctgga tgtggaacct atccttctgt    480
gtagtaaata cacctagtaa gactccatta aaggaat                                518

```

```

<210> SEQ ID NO 93
<211> LENGTH: 561
<212> TYPE: DNA
<213> ORGANISM: Mus musculus

```

```

<400> SEQUENCE: 93

```

```

aaaactctgt gtggcccagt gaccttgaac ttcccatcct tcttctccta cctctggctt    60
agtgtcttct ctttccaccc cgcacctccc cctctcaaaa aatattctta acagagggtg    120
gaaacaggtc tgaggggtgc aaagtgtgac ccaacaccag tgggggacta ggattcctct    180
ctctctaaga tagatttatt atctatctat ctatctatct atctatctat ctatctatct    240
atctatctat ctatctttat ttatttgaca aggattcctc tctctttaag atgtatttat    300
ttattttatt attttttat ttatttgaca ctgtcgcagt gttcagaccc agcaccagtt    360
ggggactagg attcctctct ctctttaaga tttatttatt tattttatta tttatttatt    420
taacactgtc acagtgttca gacacaccag aagaggggat tggaccccat tacagatggt    480
tgtgagtcac catgtgattg ctgggaatta aactcaggac ctctggaaga gcagtcagtg    540
tttttaaccg ttgggccatc t                                561

```

```

<210> SEQ ID NO 94
<211> LENGTH: 600
<212> TYPE: DNA
<213> ORGANISM: Mus musculus

```

```

<400> SEQUENCE: 94

```

```

gttttagctat cagagatgcc ccttccactc cttcattcca tcttctgaa attttttagaa    60
atgtcattta ttttgaata agagagtcca ctatggggga ctagagagat ggctcagtg    120
ttaagagcaa ggacagctct cctaaaggtc ctgaattcaa tcccagcaa ccacatggtg    180
gctcacaacc atctgtagtg ggatcccatg cctcttctcg gtgtgtctga aaagagcaat    240
gggtgactca tatacatata tacatacata catacatata tacatacata catacatata    300
tacatacata catacatata taaatcttta acaaaaaagc ctagagtcca ctatctgtcc    360
aggtecttag gctcagcgg agtgtacagc aagggagtag cctcctgggt gaggctgact    420
ttgcgcccat gccagccttt gccattttac actgtttgct tcaactagtt ttataactca    480
cccactgcac atatctccat actatccctt gtgaagtaat gataaacat gggtaatgtc    540
ttataacagt gatgtaatat acatagctat gactattatc atcatagatt tgtacttaac    600

```

```

<210> SEQ ID NO 95
<211> LENGTH: 797
<212> TYPE: DNA

```

-continued

---

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 95

```

gaattttctg tggcagaact gttctttaga ttttgtgcat ctttccaatt ttaacatfff    60
tattatftaa aaaaaataaa ttctatfttc atctcatgag cattggtgft ctgcaaacat    120
ctatgfttat atgagggtgt cagaaccttt gaaattggag ttacagatag ttgtgagcta    180
ccatgtgggt acattgtftaa gctgtaacat atttgtctgt ctaatatctg tctgtfttate    240
tatatatctg tctataatft gtcactctac tactctacta tcatactgtc tattftctatt    300
tattatctat ctatctatct atctatctat ctatctatct atctatctat ctatctatct    360
atgcatatat ctctatctat gtatctatct atgcatatat ctctatgtat ctatctatct    420
atgcatatft ctctatctat gtatcaactc ctatgtatat atctatcatt tatcatctat    480
ccatcctcta tcacatgtct attatctftc tatctatctt tacatatatt taactatctt    540
tgtatgtatg tatgtatgta tgtatgtatg tatctfttctg gatgtatgta tctatccatc    600
tttgtgtgtg tgtgtgtgtg tgtgtatata tatatcatct atcaatctat catttagcta    660
tctatatcaa tctatgcatc tatctatctc tctgtataca catatgtatg tattatctat    720
ctacctatca tctagccatc tactatcata taatatgtat atgtataata tgtatatata    780
atatgtataa tatatat                                     797

```

<210> SEQ ID NO 96

<211> LENGTH: 704

<212> TYPE: DNA

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 96

```

cccaaggcag taatatctctg gtctctggcca cctatgttag tgatctctgt tttgctctat    60
ccgagtctgg caactgacat cattfttcaat atggagacaa tggaaaccgag gaaagcagtt    120
ctftftftggg ttaggacctt taagcagcat ctaaacactca gctataaata tgaccaagta    180
aatagcfttac aattatagtc tggcagagtg catacctagc agfttatgaag ccccaagctc    240
catccccagc acagcataaa ctgggtgtag tagtggcaga cctgtfttgc tttggaaggt    300
ggaagcagaa ggctcacaaa ttgaaggaca ttatgtatgt atgtatgtat gtatgtatgt    360
atgtatgtat gtatgtatgt atgtatgtat gtatgtatft tctcgggtct ctcaatgact    420
tcttagcagt cctggaactc tgaagaccag gctggccctg aactcagaaa tttgattggt    480
attaaggtg tgcaccacca ggcccagftc taggccatc ttatccaata gctacatagft    540
gagctctaga ccattfttggg gtatattaga cactgtgtaa aagggaaaga aaaaatagac    600
aaggaagftt tatgatcaga atcactgatc aaatgtatga aacttaatft acaaacatat    660
attacctaa g cacatgtgt ctgccatctt ttctcagata caaa                               704

```

<210> SEQ ID NO 97

<211> LENGTH: 522

<212> TYPE: DNA

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 97

```

atftaggggc atftgtgaat atatactata tcttgaggat tatactagag tctgctgaca    60
gagftataac agggagagat gaaatftftc cttagtctt tgtggctgga ccgaggggga    120
gggagccctc agaaaacaac ttcaactftt gctggaaaga gaaagagtaa ataataagat    180
atftatftta aaagaatata tacatacata catacataca tacatacata catacataca    240

```

-continued

---

tacatacata catacatata tactaggata ataggaaatg acaagctaag caggaacttt	300
agaatggcaa gtggaaacag ctggcctgat agtctctacc aaggccagtg attgctcagt	360
gattaaaacc tgggaataaa tacacctctc taatattaaa agtaatgcat ggctcaacaa	420
aaaaatacat cttcacacat tatacacagc gtatgtgaca ccatgcagtg gacacatca	480
atcggaggac ctagggtggc ttgtgggaga tccaaacacc ta	522

<210> SEQ ID NO 98  
 <211> LENGTH: 540  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 98

aaagtattgc aaataattag gtcagggtg ctggcattcg gtataaggca ggggaattca	60
cagtcttgac acaggggctg gacgatggct cagcttttag gagcacttgt tgctctttca	120
gaggaccag cacctataag gtggctcaca accatcagta actccagttc caggggaatc	180
tttgacaccc cccctctctg acctctgtgg atatcaggca cacacatgga acacacctct	240
acctgaaggc aaaatactca tacacataat ttttttaaaa aatctatcta tctatctatc	300
tatctatcta tctatctatc tatctatcta tctatctatc tatctagggc cagatgtggt	360
ggtgcacaac tttaccagcc ctcaagacgt ggatctctgg gagtttgagg ccagcctggt	420
ctacatagtg agcttcaggc aaatcaggac tgcaatggtg agactctgtc tcctaaataa	480
ataaatctat ctaaaaaatt ttttaaggaa agaaaaagt tcaggccagc ctgagtgtag	540

<210> SEQ ID NO 99  
 <211> LENGTH: 540  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 99

ggacacttgt gctaataagag agtgtagccc agaaggtoat aggccagag tttctcagtg	60
ttagggaagg ctggtgtttg tgggtggtg gaaacctcac tgtaaacagt ggatggtctt	120
aaggcagcaa tgettttctc tctctctgtc catggcttgc cttgctttat gtgctccaga	180
accatcctgg catgtctcta ctcttctcca tccgtccatc cctccatccc tccatccact	240
atctatctat ctatctatct atctatctat ctatctatct atctatctat ctatctatct	300
atctcatgga ggtcagaggc cagtttggtg gagttgttcc tttctttcca gcacataact	360
tggggatcag atttagttta tcagatttgt ggcaagagcc atgtcacagg ccccccccc	420
acaccccccg ctttgtcttt ttgaagacca ggcctctca gtttgtgctt ggagtagaga	480
aatcaattat ttgaactaaa gtgattccct gtataaacag ttcacacttt aagatgaatt	540

<210> SEQ ID NO 100  
 <211> LENGTH: 600  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 100

agccaggaca tttcccttgt ttatgccaag ggctcagaaa tggcaactcc cactgcttct	60
tggatataac ccaagtcctt gaggcctcatt ctgctccacg agggacacag cagtgaggca	120
agacaggaag ggacattcac tcgttaacct gcctctttcc ctccttagct aaccttttat	180
cccaccttta ccagtatatg ttctccagta agcaacaacg atctttctat ctatctatca	240

-continued

---

```
tctatctatc tatctatcta tctatctatc tatctatcta tctatctatc tatcccatag 300
aacagctct caccttagag ttcagagttg ctttgacagg agagacacaa agagaatcct 360
gtgtgcagcc agtggcagcc aatgggctag ttccacaggt tggactcttc tctoccaatg 420
ggggtgtccc aattaggaca tgaagcctct gggagtaatg aggctgtggc ggggtggagcc 480
tagttatgac tagtgcttta ttagtgactt ttttccctta ctgtgacaaa aacacctgac 540
agaagcaact taaggagaaa gggtttactt tggcttatgg tttatgggga taacagtcac 600
```

```
<210> SEQ ID NO 101
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
```

```
<400> SEQUENCE: 101
```

```
tctgctgatt gtgccattg 20
```

```
<210> SEQ ID NO 102
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
```

```
<400> SEQUENCE: 102
```

```
attgctggt tagtgacaca 20
```

---

We claim:

1. A method of determining the alleles present in a DNA sample, the method comprising:

obtaining a DNA sample to be analyzed;

selecting a set of STR marker loci of the DNA sample to be analyzed that can be amplified together in a multiplex amplification reaction, wherein the set of STR marker loci are selected from the group consisting of: 18-3, 4-2, 6-7, 15-3, 6-4, 12-1, 5-5, X-1, Mo11-1, Mo19-2, Mo1-2, Mo7-1, Mo1-1, Mo3-2, Mo8-1, Mo2-1, Mo11-2, Mo17-2, Mo13-1, D8S1106 and D4S2408;

providing a set of oligonucleotide primer pairs, wherein each oligonucleotide primer pair includes a forward primer and a reverse primer, and each oligonucleotide primer pair in the set flanks a single locus in the set of STR marker loci, and wherein each oligonucleotide primer pair is capable of amplification of a single locus from the set of STR marker loci in a multiplex amplification reaction;

co-amplifying the set of STR marker loci in a multiplex amplification reaction, wherein the product of the multiplex amplification reaction comprises a mixture of amplified alleles from each of the co-amplified loci in the set of STR marker loci;

evaluating the products of the co-amplification reaction to determine the alleles present at each of the loci analyzed in the set of STR marker loci within the DNA sample; and

wherein the source of the DNA sample to be analyzed is at least one of a mouse and a cell line derived from a mouse.

2. The method of claim 1, wherein the mouse is selected from the group consisting of *Mus musculus musculus*, *Mus musculus domesticus*, and *Mus musculus molossinus*.

3. The method of claim 1, wherein the DNA sample to be analyzed comprises DNA derived from at least two different mouse sources.

4. The method of claim 1, wherein the DNA sample to be analyzed further comprises DNA derived from at least one other source that is a non-mouse source.

5. The method of claim 4, wherein the non-mouse source is at least one of human and African green monkey.

6. The method of claim 5, wherein the set of oligonucleotide primer pairs further comprises an oligonucleotide primer pair configured to amplify an STR allele from the non-mouse source.

7. The method of claim 1, wherein at least one oligonucleotide primer in each oligonucleotide pair comprises a detection label.

8. The method of claim 7, wherein the detection label comprises a fluorescent dye.

9. The method of claim 8, wherein evaluating the products of the co-amplification reaction comprises measuring fluorescence emission upon excitation of the detection label with light.

10. The method of claim 9, comprising using at least four oligonucleotide primer pairs, wherein at least one primer of each oligonucleotide primer pair is labeled with a detection label, and wherein at least four different detection labels are used.

11. The method of claim 1, wherein the forward primer of the oligonucleotide primer pair is SEQ. 10, the reverse primer of the oligonucleotide primer pair is SEQ. 11, and the STR marker loci is 18-3.

12. The method of claim 1, wherein the forward primer of the oligonucleotide primer pair is SEQ. 12, the reverse primer of the oligonucleotide primer pair is SEQ. 13, and the STR marker loci is 4-2.

13. The method of claim 1, wherein the forward primer of the oligonucleotide primer pair is SEQ. 14, the reverse primer of the oligonucleotide primer pair is SEQ. 15, and the STR marker loci is 6-7.

14. The method of claim 1, wherein the forward primer of the oligonucleotide primer pair is SEQ. 16, the reverse primer of the oligonucleotide primer pair is SEQ. 17, and the STR marker loci is 9-2.



57

15. The method of claim 1, wherein the forward primer of the oligonucleotide primer pair is SEQ. 18, the reverse primer of the oligonucleotide primer pair is SEQ. 19, and the STR marker loci is 15-3.

16. The method of claim 1, wherein the forward primer of the oligonucleotide primer pair is SEQ. 20, the reverse primer of the oligonucleotide primer pair is SEQ. 21, and the STR marker loci is 6-4.

17. The method of claim 1, wherein the forward primer of the oligonucleotide primer pair is SEQ. 22, the reverse primer of the oligonucleotide primer pair is SEQ. 23, and the STR marker loci is 12-1.

18. The method of claim 1, wherein the forward primer of the oligonucleotide primer pair is SEQ. 24, the reverse primer of the oligonucleotide primer pair is SEQ. 25, and the STR marker loci is 5-5.

19. The method of claim 1, wherein the forward primer of the oligonucleotide primer pair is SEQ. 26, the reverse primer of the oligonucleotide primer pair is SEQ. 27, and the STR marker loci is X-1.

20. The method of claim 1, wherein the forward primer of the oligonucleotide primer pair is SEQ. 34, the reverse primer of the oligonucleotide primer pair is SEQ. 35, and the STR marker loci is Mo11-1.

21. The method of claim 1, wherein the forward primer of the oligonucleotide primer pair is SEQ. 36, the reverse primer of the oligonucleotide primer pair is SEQ. 37, and the STR marker loci is Mo19-2.

22. The method of claim 1, wherein the forward primer of the oligonucleotide primer pair is SEQ. 38, the reverse primer of the oligonucleotide primer pair is SEQ. 39, and the STR marker loci is Mo1-2.

23. The method of claim 1, wherein the forward primer of the oligonucleotide primer pair is SEQ. 40, the reverse primer of the oligonucleotide primer pair is SEQ. 41, and the STR marker loci is Mo7-1.

24. The method of claim 1, wherein the forward primer of the oligonucleotide primer pair is SEQ. 42, the reverse primer of the oligonucleotide primer pair is SEQ. 43, and the STR marker loci is Mo1-1.

58

25. The method of claim 1, wherein the forward primer of the oligonucleotide primer pair is SEQ. 44, the reverse primer of the oligonucleotide primer pair is SEQ. 45, and the STR marker loci is Mo3-2.

26. The method of claim 1, wherein the forward primer of the oligonucleotide primer pair is SEQ. 46, the reverse primer of the oligonucleotide primer pair is SEQ. 47, and the STR marker loci is Mo8-1.

27. The method of claim 1, wherein the forward primer of the oligonucleotide primer pair is SEQ. 48, the reverse primer of the oligonucleotide primer pair is SEQ. 49, and the STR marker loci is Mo2-1.

28. The method of claim 1, wherein the forward primer of the oligonucleotide primer pair is SEQ. 50, the reverse primer of the oligonucleotide primer pair is SEQ. 51, and the STR marker loci is Mo11-2.

29. The method of claim 1, wherein the forward primer of the oligonucleotide primer pair is SEQ. 52, the reverse primer of the oligonucleotide primer pair is SEQ. 53, and the STR marker loci is Mo17-2.

30. The method of claim 1, wherein the forward primer of the oligonucleotide primer pair is SEQ. 54, the reverse primer of the oligonucleotide primer pair is SEQ. 55, and the STR marker loci is Mo13-1.

31. The method of claim 1, wherein the forward primer of the oligonucleotide primer pair is SEQ. 87, the reverse primer of the oligonucleotide primer pair is SEQ. 88, and the STR marker loci is 15-3.

32. The method of claim 1, wherein the forward primer of the oligonucleotide primer pair is SEQ. 89, the reverse primer of the oligonucleotide primer pair is SEQ. 90, and the STR marker loci is 12-1.

33. The method of claim 1, wherein the forward primer of the oligonucleotide primer pair is SEQ. 93, the reverse primer of the oligonucleotide primer pair is SEQ. 94, and the STR marker loci is X-1.

\* \* \* \* \*