

Standard for Training in Forensic Short Tandem Repeat Typing Methods using Amplification, DNA Separation and Allele Detection

*Biological Methods Subcommittee
Biology/DNA Scientific Area Committee
Organization of Scientific Area Committees (OSAC) for Forensic Science*



OSAC Proposed Standard

Standard for Training in Forensic Short Tandem Repeat Typing Methods using Amplification, DNA Separation and Allele Detection

Prepared by
Biological Methods Subcommittee
Version: 1.0
2018

Disclaimer:

This document has been developed by the Biological Methods Subcommittee of the Organization of Scientific Area Committees (OSAC) for Forensic Science through a consensus process and is *proposed* for further development through a Standard Developing Organization (SDO). This document is being made available so that the forensic science community and interested parties can consider the recommendations of the OSAC pertaining to applicable forensic science practices. The document was developed with input from experts in a broad array of forensic science disciplines as well as scientific research, measurement science, statistics, law, and policy.

This document has not been published by an SDO. Its contents are subject to change during the standards development process. All interested groups or individuals are strongly encouraged to submit comments on this proposed document during the open comment period administered by the Academy Standards Board (www.asbstandardsboard.org).



Foreword

This standard defines the minimum requirements that shall be met in a forensic DNA analyst training program for short tandem repeat typing methods using amplification, DNA separation, and detection. The aim is to provide a framework for quality training that will result in consistency in the forensic DNA community.

This standard was revised, prepared and finalized as a standard by the DNA Consensus Body of the AAFS Standards Board (ASB). The initial draft document was developed by the Biological Methods Subcommittee of the Organization of Scientific Area Committees. All hyperlinks and web addresses shown in this document are current as of the publication date of this standard.

Keywords: Training, PCR, amplification, STR typing, DNA separation, capillary electrophoresis, DNA detection

Table of Contents

1 Scope	1
2 Normative References	1
3 Terms and Definitions	1
4 Requirements	3
4.1 Knowledge-based training	3
4.2 Practical training	5
4.3 Competency testing	6
5 Conformance	7
Annex A	8

DRAFT

1 Scope

This standard provides the requirements of a forensic DNA laboratory's training program in forensic Short Tandem Repeat typing methods using amplification, DNA separation and allele detection.

2 Normative References

ASB Standard 022 - *Standard for Forensic DNA Analysis Training Programs*¹

3 Terms and Definitions

For purposes of this document, the following definitions apply.

3.1

Allele

One of two or more versions of a genetic sequence at a particular location in the genome.

3.2

Amplification

An increase in the number of copies of a specific DNA fragment; can be *in vivo* or *in vitro*. In forensic DNA testing laboratories, this refers to the use of the PCR technique to produce many more copies of specific genetic loci from samples of known and unknown origin for the purpose of generating DNA profiles for comparison.

3.3

Analytical threshold

1) The minimum height requirement at and above which detected peaks on a STR DNA profile electropherogram can be reliably distinguished from background noise; peaks above this threshold are generally not considered noise and are either artifacts or true alleles. 2) An acceptable "Relative Fluorescence Units" (RFU) level determined to be appropriate for use in the PCR/STR DNA typing process; a minimum threshold for data comparison is identified by the specific forensic laboratory doing the testing through independent validation studies.

3.4

Artifact

A non-allelic product of the amplification process (e.g., stutter, non-templated nucleotide addition, or other non-specific product), an anomaly of the detection process (e.g., pull-up or spike), or a by-product of primer synthesis (e.g., "dye blob") that may be observed on an electropherogram; some artifacts may complicate the interpretation of DNA profiles when they cannot be distinguished from the actual allele(s) from a particular sample.

¹ American Academy of Forensic Sciences Standards Board, 4200 Wisconsin Avenue, NW, Suite 106-310, Washington, DC 20016-2143 asb@aafs.org

3.5

Bin

Allele designations corresponding to the window of fragment sizes for each allele, determined by empirical testing.

3.6

Capillary electrophoresis

An electrophoretic technique for separating DNA molecules by their relative size based on migration through a narrow glass capillary tube filled with a liquid polymer.

3.7

Contamination

Exogenous DNA or other biological material in a DNA sample, PCR reaction, or item of evidence; the exogenous DNA or biological material could be present before the sample is collected, or introduced during collection or testing of the sample.

3.8

Electrophoresis

A technique used in laboratories to separate macromolecules based on size. Negatively charged molecules (e.g. proteins, DNA and RNA) migrate towards a positively charged pole through a sieving matrix, which permits a size-dependent separation.

3.9

Inhibitor

As related to the polymerase chain reaction (PCR), any substance that interferes with or prevents the synthesis of DNA during the amplification process.

3.10

Locus (plural loci)

A unique physical location of a gene (or specific sequence of DNA) on a chromosome

3.11

Polymerase chain reaction (PCR)

An enzymatic process by which a specific region of DNA is replicated during repetitive cycles that consist of the following: denaturation of the template; annealing of primers to complementary sequences at an empirically determined temperature; and extension of the bound primers by a DNA polymerase. The goal of the PCR process is to generate many copies (termed products or amplicons) of a specific region of DNA for further analysis.

3.12

Short tandem repeats (STR)

Multiple copies of an identical (or similar) DNA sequence arranged in direct succession where the repeat sequence unit is 2 base pairs (bp) to 6 bp in length; because STRs generally occur in the DNA outside of the constraints (*i.e.*, selective pressure) of genes, the number of repeat units can vary among individuals.

3.13

Spectral calibration

An examination of the contribution of overlap in the emission spectrum of fluorescent dyes used for a specific DNA test on a capillary electrophoresis instrument; permits the color deconvolution

necessary for multi-color STR typing or sequencing to be performed; a poor spectral calibration may cause artifact peaks or inaccurate peak height determinations.

3.14

Stochastic

1) Chance, or random variation 2) in DNA testing, refers to random sampling error from extracts containing low levels of DNA and/or random variation in selection of alleles amplified at a particular locus.

4 Requirements

4.1 Knowledge-based training

The laboratory's training program shall provide the trainee with an understanding of the fundamental principles of the theory behind PCR amplification, DNA separation, and allele detection methods, the function of the reagents and other components used in each method, the limitations of each method, and the laboratory's own STR typing guidelines

4.1.1. At a minimum, the knowledge-based portion of the training program shall require review of the following:

- a) The laboratory's protocols for PCR amplification, DNA separation, and allele detection
- b) The laboratory's applicable validation studies
- c) Literature used to support validation of the methods
- d) Applicable literature as assigned by the trainer

4.1.2 At a minimum, the knowledge-based portion of the training program shall cover the following topics:

NOTE: Knowledge of historical methods is intended to provide an educated perspective on current methods. In-depth understanding of these methods may not be required for successful training.

- a) STRs in forensic DNA analysis
 - i. History
 - ii. Structure and nomenclature
 - iii. Methods of analysis
 - iv. STR typing systems (e.g., commercially produced kits)
 - v. Core STR loci (e.g., CODIS)
- b) Polymerase chain reaction

- i. History
 - ii. Biochemical principles
 - a. Hot-start PCR
 - b. Multiplex PCR
 - iii. Function of reagents
 - iv. Specificity, fidelity, and optimization
 - v. Limitations
 - vi. PCR inhibitors
 - vii. Stochastic events
 - viii. Amplification artifacts
 - ix. Contamination and quality control
- c) DNA separation
- i. Theory of electrophoresis
 - ii. Capillary electrophoresis advantages and disadvantages
 - iii. Function of reagents
 - iv. Electrokinetic injection
 - v. DNA sieving
 - vi. Sample preparation
 - vii. Electrophoresis artifacts
 - viii. Contamination and quality control
- d) DNA detection
- i. History of DNA detection methods
 - ii. Fluorescent dye detection
 - a. Excitation

- b. Emission
 - iii. Dye-labeling of PCR primers
 - iv. Computer software programs for DNA detection
 - v. Multicomponent analysis/spectral calibration
 - vi. Analytical threshold
 - vii. Fragment sizing and allele calling
 - viii. Virtual bins
- e) Instrumentation and reagents
 - i. Thermal cycling instruments and parameters
 - ii. DNA separation and detection instruments and parameters
 - iii. Software parameters associated with instruments
 - iv. Maintenance and calibration
 - v. Storage of STR typing kit and DNA separation reagents
- f) Troubleshooting
 - i. Thermal cycling errors (e.g., ramping, temperature control)
 - ii. DNA detection errors (e.g., spectral calibration failure, resolution failure)
 - iii. General equipment failure

4.2 Practical training

The laboratory's training program shall provide the trainee with sufficient practical instruction for the trainee to obtain the skills for performing the STR PCR amplification, DNA separation, and allele detection protocols used by the laboratory.

- 4.2.1 At a minimum, the practical portion of the training program shall include exercises representative of the range, type, and complexity of routine casework or database samples processed by the laboratory. These include:
- a) STR PCR amplification, DNA separation, and allele detection methods to be utilized by the trainee
 - b) Documentation of the process

- 4.2.2 Practical exercises representative of the range, type, and complexity of routine casework or database samples processed by the laboratory. These shall include
- a) STR PCR amplification, DNA separation, and allele detection methods to be utilized by the trainee
 - b) Documentation of the process
 - c) The number and quality of samples processed by the trainee shall be appropriate to demonstrate the ability to follow the laboratory's STR PCR amplification, DNA separation, and allele detection protocol(s) and to produce reliable and accurate results.

4.3 Competency testing

The laboratory's training program shall include knowledge-based and practical competency testing in the application of STR PCR amplification, DNA separation, and allele detection protocols used by the laboratory. The format of the test(s) shall meet section 4.3 of ASB 022.

4.3.1 Knowledge-based competency

The trainee shall successfully complete a knowledge-based test covering the critical information obtained during the training of STR typing methods using PCR amplification, DNA separation, and allele detection methods. The test(s) shall cover, at a minimum:

- a) The theoretical and scientific bases of STR PCR amplification, DNA separation, and detection
- b) The function of the reagents and other components used in each method
- c) The proper application of each method
- d) The quality control steps pertaining to PCR amplification, DNA separation, and allele detection
- e) The laboratory's analytical procedures pertaining to PCR amplification, DNA separation, and detection

4.3.2 Practical competency

The trainee shall successfully complete a practical test covering each of the PCR amplification, DNA separation and detection protocol(s) for which he or she will be independently authorized. The trainee shall be able to satisfactorily perform the following, as applicable:

- a) Properly and accurately execute the analytical procedures related to PCR amplification, DNA separation and detection

- b) Apply the laboratory's analytical procedures to a variety of evidentiary casework or database samples
- c) Operate equipment and instrumentation used in the laboratory
- d) Document work performed in accordance with laboratory procedures

5 Conformance

In order to demonstrate conformance with this standard, the laboratory shall meet Section 5 of the ASB 022.

DRAFT

Annex A **(informative)**

Bibliography

The following information provides a list of the literature resources that may assist the DNA technical leader in defining the breadth and scope of the materials to be reviewed by the trainee. This list is not meant to be all inclusive. The laboratory shall develop a list tailored to its specific needs. Updated references shall be added to the laboratory's list as new methods or technologies are incorporated into the laboratory's protocols.

- 1] Akane, A et al., Identification of the heme compound copurified with deoxyribonucleic acid (DNA) from bloodstains, a major inhibitor of polymerase chain reaction (PCR) amplification. *J Forensic Sci* 39: 362-372, 1994
- 2] Altria, KD, Fundamentals of capillary electrophoresis theory. In *Methods in Molecular Biology* Vol. 52: Capillary Electrophoresis, ed. K. Altria, Humana Press, pp. 3-12, 1996
- 3] Bär, W, et al., DNA recommendations - 1994 report concerning further recommendations of the DNA Commission of the ISFH regarding PCR-based polymorphisms in STR (short tandem repeat) systems, *International Journal of Legal Medicine*, 107:159-160, 1994
- 4] Bär, W, et al., DNA recommendations - Further report of the DNA Commission of the ISFH regarding the use of short tandem repeat systems, *International Journal of Legal Medicine*, 110:175-176, 1997
- 5] Barron, AE and Blanch, HW, DNA separations by slab gel and capillary electrophoresis, *Separation Purification Methods* 24:1-118, 1995
- 6] Budowle, B, et al., CODIS and PCR-based short tandem repeat loci: Law enforcement tools. *Proceedings of the Second European Symposium on Human Identification*, pp. 73-88, 1998
- 7] Buse, EL, et al., Performance evaluation of two multiplexes used in fluorescent short tandem repeat DNA analysis, *J Forensic Sci*, 48:348-357
- 8] Butler, JM, *Advanced Topics in Forensic DNA Typing: Methodology*. Elsevier, 2012
- 9] Butler, JM, *Forensic DNA Typing*. Elsevier, 2005
- 10] Butler, JM, *Fundamentals of Forensic DNA Typing*. Elsevier, 2010
- 11] Butler, JM, Short tandem repeat typing technologies used in human identity testing, *Biotechniques* 43(4), Sii-Sv, 2007
- 12] Butler, JM, Genetics and genomics of core short tandem repeat loci used in human identity testing. *J Forensic Sci*. 51: 253-265, 2006
- 13] Butler, JM, et al., The development of reduced size STR amplicons as tools for analysis of degraded DNA, *J Forensic Sci* 48(5):1054- 1064, 2003

- 14] Butler, JM, et al., Forensic DNA typing by capillary electrophoresis using the ABI Prism 310 and 3100 genetic analyzers for STR analysis. *Electrophoresis* 25: 1397-1412, 2004
- 15] Chamberlain, JS and Chamberlain, JR, Optimization of Multiplex PCRs. In *The Polymerase Chain Reaction*, ed. KB Mullis et al., Birkhauser, pp. 38-46, 1994
- 16] Carey, L and Mitnik, L, Trends in DNA forensic analysis. *Electrophoresis* 23: 1386-1397, 2002
- 17] Collins, PJ, et al., Developmental validation of a single-tube amplification of the 13 CODIS STR loci, D2S1338, D19S433, and amelogenin: The AmpFISTR Identifier PCR Amplification Kit, *J Forensic Sci* 49:1265-1277, 2004
- 18] Committee on DNA Forensic Science, National Research Council. *DNA Technology in Forensic Science*. National Academy Press, Washington, DC, 1992
- 19] Committee on DNA Forensic Science, National Research Council. *An Update: The Evaluation of Forensic DNA Evidence*. National Academy Press, Washington, DC, 1996
- 20] Committee on the Conduct of Science, National Academy of Sciences. *On being a scientist*, *Proceeding of the National Academy of Sciences* 86(23):9053-9074, December 1989
- 21] Crouse, CA, et al., Analysis and interpretation of short tandem repeat microvariants and three-banded allele patterns using multiple allele detection systems, *Journal of Forensic Sciences* 44:87-94, 1999
- 22] Delamoye, M, et al., False homozygosities at various loci revealed by discrepancies between commercial kits: implications for genetic databases. *Forensic Sci. Intern.* 143:47-52, 2004
- 23] Edwards, A, et al., DNA typing and genetic mapping with trimeric and tetrameric tandem repeats, *American Journal of Human Genetics* 49:746-756, 1991
- 24] Ellegren, H, Microsatellites: Simple sequences with complex evolution, *Nature Reviews Genetics* 5:435-445, 2004
- 25] Ensenberger, M, et al., Developmental validation of the PowerPlex Fusion 6C system, *Forensic Science International: Genetics* 21:1344-144, 2016
- 26] Fregeau, CJ and Fourney, RM, DNA typing with fluorescently tagged short tandem repeats: A sensitive and accurate approach to human identification, *BioTechniques* 15:100-119, 1993
- 27] Gilder, JR et al., Run-specific limits of detection and quantitation for STR-based DNA testing. *J. Forensic Sci.* 52: 97-101, 2007
- 28] Gill, P, Role of short tandem repeat DNA in forensic casework in the UK - Past, present, and future perspectives, *BioTechniques* 32:366-372, 2002
- 29] Gill, P, et al., Considerations from the European DNA profiling group (EDNAP) concerning STR nomenclature, *Forensic Science International*, 87:185-192, 1997
- 30] Gill, P, et al., Development of guidelines to designate alleles using an STR multiplex systems, *Forensic Science International*, 89:185-197, 1997

- 31] Giusti, AM, et al., Application of deoxyribonucleic acid (DNA) polymorphisms to the analysis of DNA recovered from sperm, *Journal of Forensic Sciences* 31:409–417, 198
- 32] Hammond, HA, et al., Evaluation of 13 short tandem repeat loci for use in personal identification applications, *American Journal of Human Genetics* 55:175–189, 1994
- 33] Hares, DR, Expanding the CODIS core loci in the United States, *Forensic Science International: Genetics*, 6:e52-e54, 2012
- 34] Hares, DR, Addendum to expanding the CODIS core loci in the United States, *Forensic Science International: Genetics*, 6:e135, 2012
- 35] Hares, DR, Selection and implementation of expanded CODIS core loci in the United States, *Forensic Science International: Genetics*, 17:33-34, 2015
- 36] Heller, C, Principles of DNA separation with capillary electrophoresis, *Electrophoresis* 22:629-643, 2001
- 37] Hill, CR, et al., Concordance study between the AmpFISTR MiniFiler PCR amplification kit and conventional STR typing kits, *J Forensic Sci* 52:870-873, 2007
- 38] Holt, CL, et al., TWGDAM validation of AmpFISTR PCR amplification kits for forensic DNA casework, *J Forensic Sci* 47:66-96, 2002
- 39] Issaq, HJ, et al., The effect of column length, applied voltage, gel type and concentration on the capillary electrophoresis separation of DNA fragments and polymerase chain reaction products, *Electrophoresis* 18:1153-1158, 1997
- 40] Jeffreys, A, et al., Hypervariable 'minisatellite' regions in human DNA. *Nature* 314 (6006):67-73, 1985
- 41] Jeffreys, AJ, et al., Positive identification of an immigration test case using human DNA fingerprints, *Nature* 317:818–819, 1985
- 42] Jeffreys, AJ, et al., Individual-specific 'fingerprints' of human DNA, *Nature* 318:577–579, 1985
- 43] Ju, J, et al., Design and synthesis of fluorescence energy transfer dye-labeled primers and their application for DNA sequencing and analysis, *Analytical Biochemistry* 231:131-140, 1995
- 44] Ju, JY et al., Energy transfer primers: A new fluorescence labeling paradigm for DNA sequencing and analysis, *Nature Medicine* 2:246-249, 1996
- 45] Kimpton, CP, et al., Automated DNA profiling employing multiplex amplification of short tandem repeat loci, *PCR Methods and Application*, 3:13-22, 1993
- 46] Kimpton, CP, et al., Validation of highly discriminating multiplex short tandem repeat amplification systems for individual identification, *Electrophoresis*, 17:1283-1293, 1996
- 47] Koreth, J, et al., Microsatellites and PCR genomic analysis, *J Pathol* 178: 239-248, 1996

- 48] Kraemer, M, et al., Developmental validation of QIAGEN Investigator 24plex QS kit and Investigator 24plex GO! Kit: two 6-dye multiplex assays for the extended CODIS core loci, *Forensic Science International: Genetics* 29:9-20, 2017
- 49] Kuhl, DPA and Caskey, CT, Trinucleotide repeats and genome variation, *Cur. Opin. Genet. Devel.* 3: 404-407, 1993
- 50] LaFountain MJ, et al., TWGDAM validation of the AmpFISTR Profiler Plus and AmpFISTR COfiler STR multiplex systems using capillary electrophoresis, *J Forensic Sci* 46:1191-1198, 2001
- 51] Lander, ES, and Budowle, B, DNA fingerprinting dispute laid to rest, *Nature* 371:735-738, 1994
- 52] Lazaruk, K, et al., Genotyping of forensic short tandem repeat (STR) systems based on sizing precision in a capillary electrophoresis instrument, *Electrophoresis* 19:86-93, 1998
- 53] Leclair, B, et al., STR DNA typing: increased sensitivity and efficient sample consumption using reduced PCR reaction volumes, *J Forensic Sci* 48(5):1001-1013, 2003
- 54] Leibelt, C, et al., Identification of a D8S1179 primer binding site mutation and the validation of a primer designed to recover null alleles, *Forensic Science International* 133:220-227, 2003
- 55] Lo, YMD, Introduction to the polymerase chain reaction. In *Methods in Molecular Medicine*, Vol. 16: Clinical Applications of PCR, Humana Press, pp. 3-10, 1998
- 56] Mannucci, A, et al., Forensic application of a rapid and quantitative DNA sex test by amplification of the X-Y homologous gene amelogenin, *International Journal of Legal Medicine* 106:190-193, 199
- 57] Mansfield, ES, et al., Analysis of multiplexed short tandem repeat (STR) systems using capillary array electrophoresis, *Electrophoresis* 19(1):101-107, 1998
- 58] Markoulatos, P, et al., Multiplex polymerase chain reaction: a practical approach. *J Clin. Lab. Anal.* 16: 47-51, 2002
- 59] Momhinweg, E, et al., D3S1358: Sequence analysis and gene frequency in a German population. *Forensic Science International* 95: 173-178, 1998
- 60] Moretti, TR, et al., Validation of short tandem repeat (STRs) for forensic usage: performance testing of fluorescent multiples STR systems and analysis of authentic and simulated forensic samples, *Journal of Forensic Sciences* 46:647-660, 2001
- 61] Moretti, TR, et al., Validation of STR typing by capillary electrophoresis, *Journal of Forensic Sciences* 46:661-676, 2001
- 62] Nakahori, Y, et al., A human X-Y homologous region encodes "amelogenin", *Genomics* 9:264-269, 1991
- 63] Prince, AM, and Andrus, L, PCR: How to kill unwanted DNA, *BioTechniques* 12: 358-360, 1992
- 64] Reynolds, R, et al., Analysis of genetic markers in forensic DNA samples using the polymerase chain reaction, *Analytical Chemistry* 63(1):2- 15, 199

- 65] Ruitberg CM, et al., STRBase: a short tandem repeat DNA database for the human identity testing community, *Nucleic Acids Research* 29 (1):320- 322, 2001
- 66] Saiki, RK, et al., Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase, *Science* 239:487–491, 1998
- 67] Saiki, RK, et al., Enzymatic amplification of beta-globin genomic sequences and restriction site analysis for diagnosis of sickle cell anemia, *Science* 230: 1350-1354, 1985
- 68] Sgueglia, JB , et al., Precision studies using the ABI Prism 3100 genetic analyzer for forensic DNA analysis, *Anal. Biol. Chem.* 376: 1247-1254, 2003
- 69] Simeon, S, et al., Discrepancies between forensic identification kits explained by a laser power supply shutdown, *Forensic Science International* 164: 72-74, 2006
- 70] Slater GW, et al, Theory of DNA electrophoresis: A look at some current challenges, *Electrophoresis* 21:3873-3887, 2000
- 71] Smith, RN, Accurate size comparison of short tandem repeat alleles amplified by PCR. *BioTechniques*, 18, 122-128
- 72] Tamariz, J, et al., The application of Ultraviolet Irradiation to exogenous sources of DNA in plasticware and water for the amplification of low copy number DNA, *J. Forensic Sci.* 51: 790-794, 2006
- 73] Urquhart, A, et al., Variation in short tandem repeat sequences – A survey of twelve microsatellite loci for use as forensic identification markers, *J. of Legal Medicine*, 107:13-20, 1994
- 74] Varsha, DNA fingerprinting in the criminal justice system: An overview, *DNA Cell Biol.* 25: 181-188, 2006
- 75] Viovy, JL and Duke, T, DNA electrophoresis in polymer solutions: Ogston sieving, reputation and constraint release, *Electrophoresis* 14:322-329
- 76] Watson, JD, and Crick, FHC, A structure for deoxyribose nucleic acid, *Nature* 1953:737-738, April 1953
- 77] Wyman, AR, and White, R, A highly polymorphic locus in human DNA, *Proceedings of the National Academy of Sciences* 77:6754–6758, 1980
- 78] Yap, EPH, et al., False-positives and Contamination in PCR. In *PCR Technology: Current Innovations*, ed. HG Griffen and AM Griffen, CRC Press, pp. 249-258, 1994
- 79] Yeung, SHI, et al., Fluorescence energy transfer-labeled primers for high-performance forensic DNA profiling, *Electrophoresis* 29:2251-2259, 2008
- 80] Dawnay N, Ogden R, Thorpe RS, Pope LC, Dawson DA, McEwing R. A forensic STR profiling system for the Eurasian badger: a framework for developing profiling systems for wildlife species. *Forensic Sci Int-Gen.* 2008;2(1):47-53. doi: 10.1016/j.fsigen.2007.08.006. PubMed PMID: ISI:000263668000007.

- 81] Maria Lucia Carneiro Vieira, Luciane Santini, Augusto Lima Diniz, and Carla de Freitas Munhoz 2016 Microsatellite markers: what they mean and why they are so useful Genetics and Molecular Biology 39, 312 – 328
- 82] Peter Galbusera, Stefan van Dongen, Erik Matthysen 2000 Cross-species amplification of microsatellite primers in passerine birds Conservation Genetics Volume 1, Issue 2, 163–168
- 83] Jones, KC, Levine, KF, Banks, JD. 2002. Characterization of 11 polymorphic tetranucleotide microsatellites for forensic applications in California elk (*Cervus elaphus canadensis*). Mol. Ecology Res. 2(4): 425-427.
- 84] Kanthaswamy, S., et al. (2009). Canine population data generated from a multiplex STR kit for use in forensic casework. Journal of Forensic Sciences, 54, 829-840.
- 85] JD Kurushima, JA Collins, JA Well, HB Ernest 2006 Development of 21 microsatellite loci for puma (*Puma concolor*) ecology and forensics Molecular Ecology Notes 6(4):1260 - 1262
- 86] Linacre et al. 2011. ISFG: Recommendations regarding the use of non-human (animal) DNA in forensic genetic investigations Forensic Science International: Genetics Volume 5, Issue 5, November 2011, Pages 501-505
- 87] Menotti-Raymond M, David VA, Stephens JC, Lyons LA, O'Brien SJ. 1997. Genetic individualization of domestic cats using feline STR loci for forensic applications. J Forensic Sci. Nov; 42(6):1039-51.
- 88] Paetkau D, Strobeck C. 1994. Microsatellite analysis of genetic variation in black bear populations. Molecular Ecology. 3(5): 489-495.
- 89] Dicks KL, Webster LM, McDowall I, Muya SM, Hopper J, O'Donoghue P. 2010 Validation studies on dinucleotide STRs for forensic identification of black rhinoceros *Diceros bicornis*. Forensic Sci Int Genet. 2017 Jan;26:e25-e27.